

Outline

Boundary layer parameterization and climate

Frédéric Hourdin

June 23, 2009

1 Introduction

2 Approaches to the parameterization of the boundary layer

- Scale decomposition
- Diffusive approaches and their limitations
- Alternatives to diffusive approaches

3 Boundary layer parameterizations in climate models

- Cumulus clouds and mass flux parametrizations
- From boundary layer to deep convection
- Tracer transport

4 Conclusion


Boundary layer parameterization

Introduction

Boundary layer in the climate system

The boundary layer :

- controls energy and water exchanges with surfaces
- drives the oceanic circulation
- is associated with a large fraction of clouds



Boundary layer parameterization

Introduction

Boundary layer in the "Earth System"


Driven by the Global Change studies, climate models are more and more complex :

CO₂ cycle, CH₄, ozone chemistry, aerosols, effect of land use


⇒ coupling between atmosphere, ocean, chemistry, vegetation ...

Leading to so-called "Earth System Models".

Boundary layer is central for most of those components.



Earth System Models




Boundary layer parameterization

Introduction

Boundary layer in the "Earth System"

Example of well identified uncertainty source in Earth-System models.

The diurnal (seasonal) cycle of plant respiration is modulated by the diurnal (seasonal) cycle of the boundary layer depth




Boundary layer parameterization

Introduction

Boundary layer in large scale models

Current climate models : horizontal mesh of 20 to 400 km.

Boundary layer processes are subgrid-scale ⇒ must be "parameterized"



Parameterizations

- describe the effect of subgrid-scale processes on large scale state variables
- through a set of approximate equations based on some internal variables
- must relate those internal variables to large scale variables (closure)
- closely linked to the numerical world.

Boundary layer parameterization

Approaches to the parameterization of the boundary layer

Scale decomposition

Outline

1 Introduction

2 Approaches to the parameterization of the boundary layer

- Scale decomposition
- Diffusive approaches and their limitations
- Alternatives to diffusive approaches

3 Boundary layer parameterizations in climate models

- Cumulus clouds and mass flux parametrizations
- From boundary layer to deep convection
- Tracer transport

4 Conclusion

Boundary layer parameterization

Approaches to the parameterization of the boundary layer

Scale decomposition

Scale decomposition of the conservation equation

Conservation equation

$\mathbf{v}$: wind field
 c : conserved quantity

Lagrangian form : $\frac{dc}{dt} = 0$

Advective form : $\frac{\partial c}{\partial t} + \mathbf{v} \cdot \text{grad} c = 0$

Flux form : $\frac{\partial \rho c}{\partial t} + \text{div}(\rho \mathbf{v} c) = 0$


Scale decomposition

$\bar{X}$: "average" or "large scale" variable
 $X' = X - \bar{X}$: turbulent fluctuation

$$\frac{\partial \bar{q}}{\partial t} + \bar{\mathbf{v}} \cdot \text{grad} \bar{q} + \frac{1}{\rho} \text{div}(\rho \mathbf{v}' c') = 0$$

Under boundary layer approximations ($\partial/\partial x \ll \partial/\partial z$):

$$\frac{\partial c}{\partial t} + \mathbf{v} \cdot \text{grad } c = S_c - \frac{1}{\rho} \frac{\partial}{\partial z} \overline{w'c'}$$



$\mathbf{v}$ and c are now the large scale variables.

c : θ , u , v , water (vapor and others), chemical compounds ...

Diffusive or local formulations for the PBL

$$\overline{w'c'} = -K_z \frac{\partial c}{\partial z} \quad \longrightarrow \quad \frac{\partial c}{\partial t} = \frac{\partial}{\partial z} \left(K_z \frac{\partial c}{\partial z} \right)$$

- Analogy with molecular viscosity (Brownian motion $\leftrightarrow$ turbulence)
- Down-gradient fluxes.
- Turbulence acts as a "mixing"

Outline

- 1 Introduction
- 2 Approaches to the parameterization of the boundary layer
 - Scale decomposition
 - Diffusive approaches and their limitations
 - Alternatives to diffusive approaches
- 3 Boundary layer parameterizations in climate models
 - Cumulus clouds and mass flux parametrizations
 - From boundary layer to deep convection
 - Tracer transport
- 4 Conclusion

Turbulent diffusivity K_z

- Prandtl (1925) mixing length: $K_z = l|\overline{w'}|$ or $K_z = l^2 \frac{\partial |\mathbf{v}|}{\partial z}$
- Accounting for static stability (Ex. Louis 1979)

$$K_z = f(Ri) l^2 \left| \frac{\partial \mathbf{v}}{\partial z} \right|, \quad \text{with } Ri = \frac{g}{\theta} \frac{\partial \theta}{\left(\frac{\partial \mathbf{v}}{\partial z} \right)^2} \quad (1)$$

- Turbulent kinetic energy $\overline{w'^2} \simeq e = \frac{1}{2} [\overline{u'^2} + \overline{v'^2} + \overline{w'^2}]$

$$\frac{\partial e}{\partial t} = -\overline{w'u'} \frac{\partial u}{\partial z} - \overline{w'v'} \frac{\partial v}{\partial z} + \frac{g}{\theta} \overline{w'\theta'} - \frac{1}{\rho} \frac{\partial \overline{w'p'}}{\partial z} - \frac{\partial \overline{w'e}}{\partial z} - \epsilon$$

Ex: Mellor and Yamada $\overline{w'\phi'} = -K_\phi \frac{\partial \phi}{\partial z}$ with $K_\phi = l\sqrt{2e}S_\phi(Ri)$

Note: $\frac{\partial e}{\partial t} = 0$ (stationarity) $\implies K_z$ of form Eq. 1


Limitations of turbulent diffusion

Assumption leading to the diffusive approach:


- Turbulence as a random process
- Small scale turbulence, i.e. of size $l \ll h$ with $h = \left[\frac{1}{c} \frac{\partial c}{\partial z} \right]^{-1}$

In the planetary boundary layer

- Long range vertical transport (from the bottom to PBL top)
- Organized structures




Cloud streets on North of France (March 2009, MSG)



Radar echoes dry convective boundary layer Florida, Hiop Campaign Weckwerth et al., 1997

Limitations of turbulent diffusion

Idealized view of the dry convective boundary layer.



In the mixed layer

- Diffusive formulation

$$\overline{w'\theta'} = -K_z \frac{\partial \theta}{\partial z} = 0 \quad \text{or slightly } < 0$$

- Uniform heating by the surface

$$\frac{\partial \theta}{\partial t} \simeq \frac{\overline{w'\theta'_0}}{z_i} \quad (\text{Cste} > 0)$$

$$\overline{w'\theta'} \simeq \frac{z - z_i}{z_i} \overline{w'\theta'_0} > 0$$

Outline

Extension of diffusive formulations

- Introduction of a countergradient term

$$\overline{w'\theta'} = K_z \left[\Gamma - \frac{\partial \theta}{\partial z} \right] = 0 \quad \text{with } \Gamma \simeq 1K/km \quad (2)$$

Imposed countergradient Deardorf, 1966

Revisited by Troen & Mart, 1986, Holtzlag & Boville, 1993, based on a similarity approach.

- Non local mixing length (Bougeault)


- Higher order closures

- Mellor & Yamada 1974, hierarchy at successive orders. Complex and still local.

- Abdella & Mc Farlane, 1997, Introduce a mass flux approach to compute the 3rd order moments in a Mellor and Yamada scheme.

"Bulk" models

Constant value (or prescribed profiles) c_{ML} with discontinuities Δc at boundaries.



$$z_i \frac{\partial c_{ML}}{\partial t} = [w'c'_{z_i} - \overline{w'c'_{z_i}}] \quad (3)$$

$$\text{with } \overline{w'c'_{z_i}} = -C\Delta c \quad (4)$$

Betts, Albrecht, Wang, Suarez et al 1983

Randall et al. 1992 and Lapen and Randall, 2002: Combination of bulk models with higher order closures

Transilient matrices


Numerical formalism (after Stull 1984)

C : Air mass exchange rate matrices between model layers


For turbulent diffusions

$$\frac{\partial c_l}{\partial t} = \frac{\partial}{\partial z} \left(K_z \frac{\partial c}{\partial z} \right) \approx \frac{K_{l+1/2} (c_{l+1} - c_l) - K_{l-1/2} (c_l - c_{l-1})}{\delta z^2}$$


$$\Rightarrow C_{l,l+1} = K_{l+1/2} \frac{\delta t}{\delta z^2}, C_{l,l} = -(K_{l-1/2} + K_{l+1/2}) \frac{\delta t}{\delta z^2}, C_{l,m} = 0 \text{ for } |l-m| > 1$$




Turbulent diffusion



Asymmetric Convective Model of Pleim and Chang 1992



Mass flux schemes combined with turbulent diffusion



Separation into 2 sub-columns :

$$X = \alpha X_u + (1 - \alpha) X_d$$

ascending plume of mass flux

$$\begin{aligned} f &= \alpha \rho w_u \\ \frac{\partial f}{\partial z} &= e - d \\ \frac{\partial f c_u}{\partial z} &= e c_d - d c_u \end{aligned}$$

$$\rho \overline{w'c'} = -\rho K_z \frac{\partial c}{\partial z} + f(c_u - c_d) \quad (5)$$

Chatfield and Brost, 1987, Hourdin et. al., 2002, Siebesma, Soarez et al, 2004

Mass flux schemes combined with turbulent diffusion


Comparison with LES

Dry convective boundary layer.

Forcing : $w'\theta'_0 = 0.24 \text{ K m/s}$

geostrophic wind of 10 m/s


Thermal Plume model (Hourdin et al. 2002).



LES



SCM (1D GCM)



$$MY = -\rho K \frac{\partial c}{\partial z}$$


$$TP = f(c_u - c_d)$$

Mass flux schemes combined with turbulent diffusion

Outline


- 1 Introduction
- 2 Approaches to the parameterization of the boundary layer
 - Scale decomposition
 - Diffusive approaches and their limitations
 - Alternatives to diffusive approaches
- 3 Boundary layer parameterizations in climate models
 - Cumulus clouds and mass flux parametrisations
 - From boundary layer to deep convection
 - Tracer transport
- 4 Conclusion

Statistical cloud schemes




Probability Distribution Function of the subrid-scale water.
Cloud = fraction of the mesh where water vapor exceeds saturation.
 $\Rightarrow$ New requirement for boundary layer scheme :
give information on the subrid-scale distribution

Extension of mass flux schemes to cumulus clouds




- Computation of condensation in the ascending plume
- Additional heating by condensation within the updraft
Feedback on the mass flux f and transport
- Computation of the water PDF




1D test of the cloudy thermal plume model

Continental diurnal cycle with cumulus
ARM EUROCS case (US Oklahoma)
Rio et al. 2008




3D test of the cloudy thermal plume model

Test of the a new physical package in the LMDZ global climate model
Impact on the coverage by low clouds



Cloud cover and satellite observations



Outline


- 1 Introduction
- 2 Approaches to the parameterization of the boundary layer
 - Scale decomposition
 - Diffusive approaches and their limitations
 - Alternatives to diffusive approaches
- 3 Boundary layer parameterizations in climate models
 - Cumulus clouds and mass flux parametrisations
 - From boundary layer to deep convection
 - Tracer transport
- 4 Conclusion

Parameterization of deep convection

Classical parameterizations :

- Mass flux schemes
- Importance of cloud phase changes and rainfall
- Controlled by instability above cloud base

Example of the Emanuel (1991) scheme :




Trigerring :
 $B(LCL+40hPa) > |CIN|$
Closure :
 $M_B = f(CAPE)$
CAPE : Convective Available Potential Energy
CIN : Convective INhibition.

A systematic bias of parameterized convection

Climate models with parameterized convection tend to predict continental convection in phase with insolation, while it peaks in late afternoon in reality and in Cloud Resolving Models (mesh ≈ 1 km).


An idealized case of continental cycle with deep convection

ARM, Oklahoma, after Guichard et al. 2004




Deep convection preceded by a phase of shallow cumulus convection
Boundary layer : preconditioning and trigerring of deep convection


ARM case with the standard LMD SCM



Control of deep convection by sub-cloud processes



New approach (Grandpeix et al. 2009) :
Control of deep convection by sub-cloud processes.
By analogy with a nozzle above a wall of height h .



ALP closure

Available Lifting Energy for the convection

Scaling with w^2 .

Triggerring : $ALE > |CIN|$

Available Lifting Power for the convection


Scaling with w^3 .

Closure : $M_B = f(\text{ALP})$

New requirements for the boundary layer scheme :

give reasonable estimates of $\overline{w'^2}$ and $\overline{w'^3}$.

Statistical cloud schemes




Boundary layer parameterization

Boundary layer parameterizations in climate models

From boundary layer to deep convection

ARM case with ALP closure, thermals and wakes




Boundary layer parameterization

Boundary layer parameterizations in climate models

From boundary layer to deep convection

ARM case with ALP closure, thermals and wakes




Boundary layer parameterization

Boundary layer parameterizations in climate models

From boundary layer to deep convection

Diurnal cycle of deep convection in the 3D LMDZ GCM



Boundary layer parameterization

Boundary layer parameterizations in climate models

Tracer transport

Outline

- 1 Introduction
- 2 Approaches to the parameterization of the boundary layer
 - Scale decomposition
 - Diffusive approaches and their limitations
 - Alternatives to diffusive approaches
- 3 **Boundary layer parameterizations in climate models**
 - Cumulus clouds and mass flux parametrisations
 - From boundary layer to deep convection
 - **Tracer transport**
- 4 Conclusion


Boundary layer parameterization

Boundary layer parameterizations in climate models

Tracer transport

Boundary layer and transport of atmospheric tracers

Test of ^{222}Rn transport : emitted on continents only



Boundary layer parameterization

Boundary layer parameterizations in climate models

Tracer transport


Boundary layer and transport of atmospheric tracers

Contribution of the biosphere to the CO₂ latitudinal contrasts

Idealized seasonal cycle for surface emission (null annual mean)

GCM and transport models from the Transcom exercise

After Dargaville et al.




Boundary layer and transport of atmospheric tracers

Concluding remarks

NO_x computation at Dome C, Antarctica

MAR Regional model



- Parameterization of boundary layer processes is a key issue for climate modeling and climate change studies.
- Climate models are more and more complex but the realism of the "new components" (chemistry, vegetation, ...) highly depends on the representation of atmospheric processes in general and boundary layer in particular.
- In current climate models (and still for a while), boundary layer processes must be parameterized.
- Boundary layer schemes must be valid from equator to pole, and from dry stable atmosphere to deep convection conditions.
- The "new components" put new constraints on boundary layer schemes.
- There is a large place for improvement of boundary layer parameterization.
- The combined use of a turbulent diffusion for small scales and mass flux schemes for organized structures seems a promising way.
- A hierarchy of approaches are available to improve and evaluate boundary layer parameterizations : 1D versus LES , 3D, nudged, weather forecast and climate, etc.