

Evaluation and improvement of the cold pool parameterization in the LMDZ climate model against Large Eddy Simulations

Mamadou Lamine Thiam¹, Frédéric Hourdin¹, Jean-Yves Grandpeix¹, Catherine Rio², and Maëlle Coulon–Decorzens¹

¹LMD/IPSL/SU/CNRS

²CNRM/MeteoFrance/CNRS

Correspondence: Mamadou Lamine Thiam (lamine.thiam@lmd.ipsl.fr)

Abstract. Cold pools, formed under clouds by the evaporation of precipitation, play a central role in maintaining and organizing atmospheric convection. It is suspected that their absence in climate models may lead to significant errors in the representation of convection, such as the premature convection extinction after sunset. The introduction of a cold pool parameterization into the LMDZ climate model has significantly improved the representation of convection, in particular its diurnal cycle. Thanks to this indirect evaluation, the parameterization of cold pool was retained in the further versions of LMDZ. However, no **detail** evaluation of the representations of cold pools themselves was done so far. This work provides for the first time such an evaluation based on Large Eddy Simulation (LES). The evaluation is done both on case of radiative–convective equilibrium that allows the study of convective processes in a well-controlled and quasi-stationary framework, without the influence of large-scale dynamics, as well as on the time-dependent continental case AMMA (African Monsoon Multidisciplinary Analysis), representative of typical Sahelian conditions with local initiation during the afternoon. First, we evaluate the physical relationships underlying the cold pool model in the LES, then, in a second step, its behavior when coupled with the deep convection scheme in the single-column version of LMDZ. The analyses carried out demonstrate the relevance of the assumptions underlying the parameterization. The initial version actually captures the main characteristics of LES cold pools but also exhibits some biases. We show how substantial modifications to the cold pool scheme and a readjustment of certain free parameters helped reduce those biases significantly. The remaining flaws could be corrected by adding convective mixing through thermal plumes within the cold pools and by modeling the evolution of cold pools number density rather than imposing it.

Copyright statement. TEXT

1 Introduction

During thunderstorms, precipitation forms inside convective clouds. When it falls either to the side or below the cloud, into air that is not saturated with water vapor, a portion of this precipitation evaporates. This evaporation cools the air, making it denser and creating so-called unsaturated downdrafts. ~~Below cloud base, this cold air spreads out horizontally. When approaching the surface the descending air masses diverge horizontally. The cold air may then accumulate in pools that expand.~~ These

expanding masses of cold air are generally called "cold pools" or "wakes". The later term was retained when developing the code and will be used here in the equations for consistency. The former term will be used throughout the text of this paper. The cold pool, that spreads horizontally because its air is ~~warmer~~ colder than its environment, can also be seen as a density current (Charba, 1974; Droegemeier and Wilhelmson, 1987). Cold pools are most often associated with a gust front, capable of lifting the surrounding warm air and thus promoting the development of new convective cells. In organized propagative systems such as squall lines, convective columns are permanently generated by cold pool fronts at the front of the system (Rotunno et al., 1988; Weisman and Rotunno, 2004).

Present over both land and ocean, cold pools are generally deeper, colder, and propagate more rapidly over land. They play a key role in the self-aggregation of tropical convection (Jeevanjee and Romps, 2013), as well as in the transition between shallow and deep convection (Khairoutdinov and Randall, 2006; Böing et al., 2012).

In atmospheric Global Circulation Models (GCMs), as those used for climate change studies, convection has to be parameterized due to the coarse horizontal resolution (30 to 300 km). Simulating convective rainfall with parameterized physics is challenging (Randall et al., 2003). GCMs often underestimate rainfall rates (Kendon et al., 2012; Pantillon et al., 2015; Tan et al., 2018; Rooney et al., 2022) and produce peak precipitation over land at noon, in phase with insolation, while the maximum precipitation is generally observed in late afternoon or during night (Randall et al., 2003; Guichard et al., 2004; Stephens et al., 2010; Dirmeyer et al., 2012). Cold pools probably play a key role in this timing, by self-maintaining convection (Pantillon et al., 2015; Grant et al., 2018). Cold pools also play a role in triggering elevated convection (i.e., convection initiated ~~at a~~ certain altitude above the boundary layer), a process that should be taken into account in GCMs to improve the representation of nocturnal precipitation over the Southern Great Plains in cenbtral US (Park et al., 2024).

One of the first attempts to parameterize cold pools was proposed by Qian et al. (1998). Later on, Grandpeix and Lafore (2010) proposed an autonomous parameterization based on a population of identical circular cold pools that are cooled by the evaporation of precipitation, this cooling term being provided by the parameterization of deep convection. This new scheme was ~~further~~ coupled to the Emanuel (1991) deep convection scheme and has since become a key parameterization of the LMDZ global climate model ~~Rio and Hourdin (2008)~~ (Rio and Hourdin, 2008). Other parameterizations of cold pools were proposed more recently, either independent from the convective model as is the case in LMDZ (Rooney et al., 2022; Freitas et al., 2024) or as part of it (Suselj et al., 2019). The works of Rooney et al. (2022) and Freitas et al. (2024) introduced propagation from one model grid cell to another, a process which is not represented so far in LMDZ although it is known to be important for representing their impact on convective organization (Freitas et al., 2024).

Nevertheless, the coupling of the Emanuel (1991) parameterization of deep convection with cold pool parameterization (Grandpeix and Lafore, 2010) and with the thermal plume model of Rio and Hourdin (2008) in the LMDZ climate model significantly improved the simulation of the diurnal cycle of precipitation over land in the tropics (Rio et al., 2009), shifting its maximum from noon to mid afternoon. A further improvement was brought by the introduction of the stochastic triggering of deep convection (Rochetin et al., 2014) which made the simulated convection more intermittent. Despite this success, and the use of the cold pool model in the standard version of the LMDZ atmospheric and IPSL (Institut Pierre Simon Laplace) coupled models (Hourdin et al., 2020; Boucher et al., 2020), ~~it~~ the representation of cold pools was not evaluated ~~in detail~~ so far. The

same applies for the other existing cold pool parameterizations (Park, 2014; Del Genio et al., 2015; Rooney et al., 2022; Freitas et al., 2024), which have not been evaluated ~~in-detail~~ for their ability to simulate the cold pools themselves, but rather in terms of their effect on convection. This is explained not only by a lack of observational data but also by the fact that the internal variables of parameterizations are not directly accessible from observations.

Large Eddy Simulations (LES) are a useful complement to observations. Their fine horizontal resolution enables them to simulate explicitly turbulent and convective motions in the boundary layer (Brown et al., 2002; Siebesma et al., 2003). One advantage of LES compared to observations is that they provide full three-dimensional information. They have been used to develop and evaluate boundary layer and convection parameterizations (Rio et al., 2009; Dorrestijn et al., 2013; Strauss et al., 2019; Suselj et al., 2019; Legay et al., 2025). Suselj et al. (2019) used LES to evaluate in detail the internal variables of their unified convection (dry, shallow and deep) parameterization, for example by validating the surface fraction covered by moist updrafts. They also used LES to validate an approximation regarding the timing of when cold pools begin to invigorate convection in their unified convection scheme, and to study the effect of cold pools in the convection. Other studies have used LES to better understand cold pool processes (Feng et al., 2015; Meyer and Haerter, 2020; Lochbihler et al., 2021) and to guide the development of cold pool parameterizations (Kurowski et al., 2018). However, to our knowledge, no study has yet exploited LES to evaluate in detail the internal variables of a cold pool parameterization.

Here we propose to use LES to evaluate ~~in-detail~~ the parameterization of cold pools of LMDZ (Grandpeix and Lafore, 2010; Grandpeix et al., 2010). We first use LES to evaluate some of the fundamental relationships between large scale state variables (for LES, the horizontal average over the domain) and internal variables which are at the basis of the parameterization. We then propose improvements which are further assessed in simulations with a Single-Column-Model (SCM) version of LMDZ against LES. In such simulations, the parameterization interacts with all the other parameterizations and depend on the values of a number of free parameters. To explore the sensitivity of the results to these free parameters and retune the model after improvement of its physical content, we use a tool for automatic calibration, High-Tune-Explorer, developed recently (Couvreur et al., 2021; Hourdin et al., 2021). This tool, based on history matching, can be used to characterize the subspace of parameter values for which the model is in agreement with LES, given a series of target metrics and associated tolerances to error (Couvreur et al., 2021). It is used here to explore the sensitivity of the agreement between SCM simulations and LES to the model free parameters.

The paper starts by presenting in section 1 the tools used: the LMDZ model, the cold pool parameterization by Grandpeix and Lafore (2010) (referred to as the GL10 hereafter), and the LES used for evaluation. The presentation of the tuning tool (largely published) and the setup of its use is let to an appendix to concentrate on model physics and improvement in the core of the paper. In section 2, we detail the cold pool sampling in LES, designed to assess the physical laws internal to the cold pool parameterization and its coupling with deep convection. Section 3 is devoted to a comparison of cold pool model variables simulated by LMDZ in SCM mode and those calculated in LES, in order to identify the model's limitations. These results will then be discussed, and proposed improvements will be ~~detailed~~ described in section 4. Finally, we conclude with a synthesis and discussion of prospects in section 5.

2 Tools and methods

2.1 LMDZ and its single-column version

LMDZ is the General Circulation Model (GCM) used in this work. Developed in the 1970s at Laboratoire de Meteorologie
 95 Dynamique (Sadourny, 1984; Hourdin et al., 2006). It is based on simplified Navier-Stokes equations for fluid mechanics, as
 well as transport equations. It represents the second generation (Hourdin et al., 2013) of a climate model initially described
 by Sadourny and Laval (1984). LMDZ is the atmospheric component of the IPSL coupled model. The latter is one of around
 twenty coupled models taking part regularly in the international model intercomparison project (CMIP), the results of which
 are used in IPCC (Intergovernmental Panel on Climate Change) reports. We use here the LMDZ6A configuration of LMDZ
 100 designed for CMIP6 and described by Hourdin et al. (2020).

LMDZ consists of two main parts, from a physical, mathematical and computational point of view. The first part, called “the
 dynamics”, concerns the numerical resolution of the atmospheric general circulation equations. This component manages hor-
 izontal exchanges between the model’s grid cells. The second part, called “physics”, calculates the impact of radiation, small-
 scale processes (subgrid) and phase changes of water on dynamic variables via “physical parameterizations”. This “physical”
 105 part is made up of juxtaposed atmospheric columns, which do not interact with each other. Within each column, the variables
 are assumed to be statistically homogeneous in the horizontal plane.

The SCM version of LMDZ is built by extracting an atmospheric column from the GCM, incorporating all subgrid-scale pa-
 rameterizations, and running it in a large-scale constrained environment. This approach has become central in the development
 and tuning of parameterizations of convection and associated clouds in several climate modeling groups (Zhang et al., 2016;
 110 Gettelman et al., 2019). Parameterizations are often developed and evaluated within this single-column framework by compar-
 ing them with LES of the same atmospheric column. The SCM/LES approach was promoted in particular by GCSS (GEWEX
 Cloud Systems Study), a program aimed at improving the parameterization of cloud systems in climate models (Krueger et al.,
 2016). A major advantage of the SCM is its low computational cost, which allows a large number of simulations, even on a
 laptop, making it particularly useful in the development phase, where extensive testing is required.

115 2.2 Convective parameterizations in LMDZ

The role of convective parameterizations is to provide sources of heating Q_1 and moistening Q_2 (following notations first
 introduced by Yanai et al., 1973) to the conservation equations of potential temperature θ and specific humidity q_v :

$$C_p \frac{D\theta}{Dt} = \frac{\theta}{T} [Q_R + (L_v + f_g L_f)(c - e)] - C_p \frac{1}{\rho} \frac{\partial \overline{\rho w' \theta'}}{\partial z} = \frac{\theta}{T} [Q_R + Q_1], \quad (1)$$

$$\frac{Dq_v}{Dt} = e - c - \frac{1}{\rho} \frac{\partial \overline{\rho w' q'_v}}{\partial z} = -Q_2 / L_v, \quad (2)$$

120 where C_p is the heat capacity of dry air, Q_R is the radiative heating, c and e are condensation and evaporation rates, f_g is
 the condensate ice fraction, L_v is the latent heat of vaporization, L_f the latent heat of fusion and $-\partial_z \overline{\rho w' \phi' / \rho}$ (with $\phi = \theta$
 or q_v) is the vertical convergence of the Reynold turbulent flux of ϕ accounting for the effect of the subgrid-scale turbulent

or convective motions on the explicitly resolved large scale flow. Note that the θ/T factor on the right hand side of the first equation was missing in the equation of Grandpeix and Lafore (2010) but not in the code of the parametrization. The convective parameterizations also often provide a source term Q_3 for momentum but it is not involved in the coupling with cold pools described here.

Note that the equations above are simplified assuming that the ice fraction f_g is unchanged by evaporation and condensation. Note also that Q_2 is a sink of humidity expressed conventionally as a heating term with constant L_v .

The parameterization of turbulence, convection and clouds in LMDZ is based on a multi-scale, or object view.

130 2.2.1 The small scale turbulence

The small scale turbulence, mainly active near the surface, is accounted for following Yamada (1983) scheme, with an eddy diffusive approach in which the eddy diffusivity relies on a prognostic equation for the turbulent kinetic energy.

2.2.2 Shallow convection

Shallow convection (dry or cloudy) is handled in LMDZ with the so-called “thermal plume model”, a mass flux scheme which was developed to account for non local vertical transport by organised thermal plumes, cells or rolls in the convective boundary layer (Hourdin et al., 2002; Rio and Hourdin, 2008). The combination of an eddy diffusion with a mass flux scheme for the representation of turbulent transport in the convective boundary layer, first proposed by Chatfield and Brost (1987), has since be popularized as the EDMF (Eddy Diffusion Mass Flux) approach. In the thermal plume model, more specifically, the population of convective structures within a grid cell are summarized into a unique effective ascending plume, with surface fraction α_{th} and a unique ascending mass flux $f_{th} = \rho\alpha_{th}w_{th}$, compensated by a mass flux $-f_{th}$ in a fraction $1 - \alpha_{th}$ of the grid cell. The sources Q_1^{th} and Q_2^{th} only contain the vertical convergence of the mass flux transport ($\overline{\rho w' \phi'} = f_{th}(\phi_{th} - \phi)$ where ϕ_{th} is the value of variable ϕ within the thermal plume), the part coming from the condensation or evaporation being treated in the so-called large-scale condensation scheme.

2.2.3 The large-scale condensation scheme

145 The large-scale condensation scheme is used to predict the cloud fraction except for deep convection, based on a probability distribution function (PDF) of the total water within the horizontal grid cell (giving the cloud fraction as the part of the grid cell with humidity above saturation). This statistical cloud scheme provides to first order: $Q_1^{lsc} = (L_v + f_g L_f)(c - e)$ and $Q_2^{lsc} = -L_v(c - e)$. For shallow cumulus or strato-cumulus, cloud condensation is thus treated outside the thermal plume scheme. Both schemes are however coupled together when the thermal plume is active within a grid cell. In this case, the subgrid water PDF is prescribed as a bimodal function, with one mode corresponding to the thermal plume and the other one to its environment. This coupling led to a strong improvement in the representation of cumulus and stratocumulus clouds (Jam et al., 2013; Hourdin et al., 2019).

2.2.4 Deep convection

Deep convection is represented with the Emanuel (1991) mass flux scheme modified by Grandpeix et al. (2004). Its fundamental principles are retained, with a representation of a population of cumulonimbus clouds as a collection of saturated updrafts and downdrafts together with an unsaturated downdraft (mass flux scheme). This parameterization simultaneously represents transport, condensation, cloud formation, and precipitation.

The core of the cumulonimbus is represented as an undiluted updraft that does not entrain air laterally above cloud base, but is gradually “eroded” while rising. This updraft is assumed to be fast enough to carry the liquid or solid water condensed within it. Following the Episodic Mixing and Buoyancy Sorting approach, a population of diluted ascending or descending ~~air masses, saturated,~~ saturated air masses are created by mixing a fraction of air shed from the adiabatic ascent with ambient air according to an imposed PDF. ~~‡~~ The spectrum of mixing fraction is divided into bins defining a population of air parcels that are lofted to their neutral buoyancy level (layer), thus creating a matrix in which each term is an exchange of air between two layers of the model. Before forming these mixtures, the water in excess in the air shed from the adiabatic updraft is precipitated, and then again, the excess water is precipitated before the diluted updraft is detrained into the environment (there is no further precipitation from diluted downdrafts). Finally, the unsaturated ~~downdrafts receive~~ downdraft receives all the rain formed during shedding or detraining in the environment. Some of these descents take place outside the clouds, in air that is not saturated with moisture, allowing them to evaporate. Below the base of the clouds, all of the precipitation is outside the clouds. The evaporation of very large amounts of rain beneath the cloud base intensifies the downdrafts, which gives rise to cold pools.

In practice, the convective tendencies Q_1 and Q_2 are separated into two parts, “saturated” and “unsaturated”. The saturated tendencies Q_1^{sat} and Q_2^{sat} take into account adiabatic updraft, diluted updrafts and downdrafts, and a downward flux in the environment compensating all these mass fluxes. The unsaturated tendencies Q_1^{unsat} and Q_2^{unsat} take into account unsaturated downdraft as well as compensating ascent.

175 2.3 The cold pool model

The cold pool model has been entirely described in GL10. Here, we introduce the main equations and internal variables of the parameterization relevant for the rest of the paper.

The cold pool model represents a population of circular cold pools (or wakes) over an infinite plane, consistently with the way the Reynolds decomposition is used to separate, in GCMs, the 3D dynamical core from the 1D world of parameterizations of subgrid transport, assuming that the statistics of the turbulent or convective motions responsible for this transport are horizontally homogeneous on an infinite plane. **Citer le Rio et Hourdin de l’encyclopédie**

The centers of those cold pools are randomly distributed with a uniform number density D_{wk} (number of cold pools per unit area) assuming no overlap between two of them. Due to the complex life cycle of cold pools (including birth, death, collisions and merging), calculating their number density requires another parameterization. So far, the value of the cold pool number




Figure 1. Conceptual diagram of the cold pool model and its coupling with convection in LMDZ climate model. (wk) denotes the cold-pool regions, characterized by a lower temperature (θ_{wk}), while (ex) represents the environment of cold pools, with a higher temperature (θ_{ex}). W_{wk} and W_{ex} denote the vertical velocities in (wk) and (ex), respectively. $\delta\theta$ and δW represent the differences in potential temperature and vertical velocity between regions (wk) and (ex). p_{wk} represents the pressure corresponding to the height of the top of cold pool, defined as the altitude where θ_{wk} becomes equal to θ_{ex} ($\delta\theta = 0$). p_{upper} represents the level at which all cold-pool model variables vanish. C_* represents the spreading speed of cold pool. W_b corresponds to the vertical velocity at cloud base. e_{wk} represents the entrainment above p_{wk} . Finally, ALE and ALP represent, respectively, the lifting energies and powers provided by thermals (ALE_{th} , ALP_{th}) and cold pools (ALE_{wk} , ALP_{wk}).

185 density is imposed. In LMDZ6A, it is fixed to a different value over ocean (10 cold pools over 100 km×100 km) and over land (8 cold pools over 1000 km×1000 km).

An important simplification of the model consists in assuming that, within a given column of the model and at a given time step, all the wakes have the same height, radius, and vertical profiles of thermodynamic variables which is equivalent to say that we are representing the population of cold pools through a mean effective cold pool. All those variables however evolve
190 in time according to the cold pool physics, as ~~detailed-explained~~ below.

In practice, the model divides the space into two parts, the interior (w_k) and exterior (or environment) of cold pools (ex) respectively. For temperature, humidity and vertical velocity, we introduce a wake anomaly $\delta X = X_{w_k} - X_{ex}$ defined as the difference of its mean values in the two subdomains and \bar{X} the average over the horizontal domain.

The main driver of the parameterization is the cooling associated with convective unsaturated downdrafts. All this cooling,
195 provided by the deep convection scheme, is applied to the interior of the cold pools thus creating a contrast in temperature δT between the interior and exterior of the pool (see Fig. 1). Once the cold pools are initiated (which requires a prior activation of deep convection), they enter a positive feedback loop with deep convection. Cold pools reinforce deep convection both because the ~~undiluted ascent, at the core of the parameterization of deep convection, is~~ saturated drafts are fed by air coming from the environment of the cold pool-pools (warmer near the surface than the average grid cell), and because of ~~a~~ the lifting energy
200 provided by the cold pool spreading as detailed in the following section. The temperature difference between the cold pools and their environment is controlled by competing effects: it increases due to cooling by convective downdrafts and decreases due to cold pools-pool spreading and gravity wave damping. When deep convection stops, this temperature ~~contrasts-decreases~~ until-switching-off contrast decreases until the cold pool parameterization is switched off based on a combination of thresholds. ~~Details of the scheme are given hereafter.~~

205 The top height of the cold pool is defined as the altitude h_{w_k} (and associated pressure p_{w_k}) where the temperature difference between (w_k) and (ex) becomes zero. Below this level cold pools are cooler than their exterior. The boundary between the cold pool and the environment is considered to be infinitely thin, and at each point on this boundary, the cold pool spreads at a rate C . C is considered to be a random variable whose mean C_* will give the rate at which the cold pool spreads. In the GL10 model, C_* scales with the square root of the potential energy available in the cold pools, i.e the energy associated with their
210 collapse, $WAPE$ (Wake Available Potential Energy), given by:

$$WAPE = g \int \frac{\delta \rho}{\bar{\rho}} = -g \int_0^{h_{w_k}} \frac{\delta \theta_v}{\theta_v} dz. \quad (3)$$

So that:

$$C_* = k \sqrt{2WAPE}, \quad (4)$$

where ρ is the air density; θ_v is the virtual potential temperature.

215 Coefficient k in equation (4) should take a value between 0 and 1. Based on 3D CRM (Cloud Resolving Models) simulations, Lafore (2000) estimated this coefficient to 0.33 in the case of a linear structure such as squall line. This is the value retained in LMDZ6A.

When cold pools appear in a grid cell where they were not present before, their fractional cover $\sigma_{\text{wk}} = \pi r^2 D_{\text{wk}}$ (where r is the cold pool radius) is set arbitrarily to 2%, corresponding to an initial radius of about 2.5 km over ocean and 30 km over land with the values chosen for the wake number density in LMDZ6A. The cold pool radius then ~~grow~~grows with rate $\dot{r} = C_*$ so that the time evolution of the fractional ~~surface~~area reads:

$$\partial_t \sigma_{\text{wk}} = 2\pi r C_* D_{\text{wk}} = 2C_* \sqrt{\pi D_{\text{wk}} \sigma_{\text{wk}}}, \quad (5)$$

This surface fraction increases over time. It is limited to 40% of the domain. When this threshold is reached the growth of the cold pools is stopped and the radius remains constant.

It is assumed that below the top of cold pool (p_{wk}), the vertical velocity profile associated with cold ~~pool~~pools results solely from the spreading at the surface, without lateral entrainment ($e_{\text{wk}} = 0$) or detrainment ($d_{\text{wk}} = 0$) between the cold pool and its environment. Above this level, the subsidence induces a lateral convergence of air feeding the cold pool which can be reinforced by additional reevaporation of rainfall below stratiform clouds. The shape of the vertical profile of the ~~velocity~~vertical p-velocity difference $\delta\omega$ (where ω is the vertical ~~velocity~~p-velocity in Pa/s, while we will note w the vertical velocity in m/s) between the cold pool region and its environment is imposed as a piecewise linear function of pressure: $\delta\omega$ increases linearly from zero at the surface up to a maximum value at p_{wk} and then decreases linearly between p_{wk} and a minimum pressure p_{upper} corresponding to the upper bound of the cold pool model. The vertical subsidence which thus increases downward between p_{upper} and p_{wk} is fed by lateral entrainment e_{wk} without detrainment so that $e_{\text{wk}} = \sigma_{\text{wk}}(1 - \sigma_{\text{wk}})\partial_p \delta\omega + \partial_t \sigma_{\text{wk}}$. This lateral entrainment accounts for the horizontal component of the meso-scale circulation known to entrain dry and warm (in terms of potential temperature) air from low- or mid- tropospheric air into the cold pool (see Fig. 1), in turn reducing the WAPE.

At p_{upper} , the top of the cold pool model, ~~$\delta e_x - \delta X$~~ cancels for all cold pool ~~state variables~~variables X . In GL10 model and in LMDZ6A, p_{upper} was set to 600 hPa and there was also a nonzero velocity difference ($\delta\omega^{cv}$) at p_{upper} , accounting for the difference of the convective mass fluxes between (wk) and (ex).

In the version used here, $\delta\omega^{cv} = 0$ above this level while it was not the case in GL10. We realized that the GL10 version was introducing a double counting of the vertical mass flux of ~~saturated~~unsaturated downdraft at p_{upper} . This modification has however little effect on the parameterization results since the mass flux associated with downdrafts is small at p_{upper} (result not shown).

As already said, it is the cooling associated with the re-evaporation of rain in unsaturated downdrafts that is the primary driver of cold pool development. This process is reflected in the model by assigning the heating term Q_1^{unsat} computed by the convective scheme to the interior of cold pools, while Q_1^{sat} acts on their environment. Consistent with this splitting, we assume that the saturated part of the convective scheme sees the profiles outside the cold pools, and the unsaturated downdrafts their interior. We further assume that thermal plumes are only active in the fraction of the horizontal surface located outside the cold pools. Thus the mass flux transport scheme is applied to any variable outside the cold pool $\phi_{\text{ex}} = \phi - \sigma_{\text{wk}}\delta\phi$, and finally $(\theta/T)Q_1^{\text{th}} = -C_p(1 - \sigma_{\text{wk}})\partial_z \overline{\rho w' \theta'_{\text{ex}}}$ and $Q_2^{\text{th}} = L_v(1 - \sigma_{\text{wk}})\partial_z \overline{\rho w' q'_{v\text{ex}}}$. The thermal plume model therefore induces a differential tendency that is opposite of the average tendency restricted to the environment. Ultimately, the contrast in convective

tendencies (shallow and deep) between the cold pools and their environment reads:

$$\begin{cases} \delta Q_1^{\text{cv}} = \frac{Q_1^{\text{unsat}}}{\sigma_{\text{wk}}} - \frac{Q_1^{\text{sat}}}{1-\sigma_{\text{wk}}} - \frac{Q_1^{\text{th}}}{1-\sigma_{\text{wk}}}, \\ \delta Q_2^{\text{cv}} = \frac{Q_2^{\text{unsat}}}{\sigma_{\text{wk}}} - \frac{Q_2^{\text{sat}}}{1-\sigma_{\text{wk}}} - \frac{Q_2^{\text{th}}}{1-\sigma_{\text{wk}}}. \end{cases} \quad (6)$$

It is the terms δQ_1^{cv} and δQ_2^{cv} that drive the time evolution of $\delta\theta$ and δq_v given by:

$$255 \quad \begin{cases} \partial_t \delta\theta = \frac{\theta}{T} \left(\frac{\delta Q_1^{\text{cv}} + \delta Q_1^{\text{wk}}}{C_p} \right) - \bar{\omega} \partial_p \delta\theta - \frac{K_{gw}}{\tau_{gw}} \delta\theta, \\ \partial_t \delta q_v = - \frac{\delta Q_2^{\text{cv}} + \delta Q_2^{\text{wk}}}{L_v} - \bar{\omega} \partial_p \delta q_v. \end{cases} \quad (7)$$

This time evolution also includes a differential heating and moistening induced by the cold pools themselves of the air inside and outside the cold pools, under the effect of lateral air entrainment from the environment above p_{wk} , subsidence inside the cold pools, and compensating ascent in the environment:

$$\begin{cases} \frac{\theta}{T} \frac{\delta Q_1^{\text{wk}}}{C_p} = - \frac{e_{\text{wk}}}{\sigma_{\text{wk}}} \delta\theta - \delta\omega \partial_p \bar{\theta} - (1 - 2\sigma_{\text{wk}}) \delta\omega \partial_p \delta\theta, \\ - \frac{\delta Q_2^{\text{wk}}}{L_v} = - \frac{e_{\text{wk}}}{\sigma_{\text{wk}}} \delta q_v - \delta\omega \partial_p \bar{q}_v - (1 - 2\sigma_{\text{wk}}) \delta\omega \partial_p \delta q_v. \end{cases} \quad (8)$$

260 The terms $-\bar{\omega} \partial_p \delta$ in (7) partially compensate for the fact that the contrasts δ are not transported by the dynamics until now. We therefore take into account, in the parameterizations, the vertical part of large-scale advection to partially compensate for this deficiency.

Finally, the last term, present only in the θ part of (7), corresponds to the reduction in temperature contrasts by gravity waves with a coefficient specified as the ratio of an efficiency K_{gw} to a characteristic time

$$265 \quad \tau_{gw} = \frac{\sqrt{\sqrt{\sigma_{\text{wk}}} - (1 - \sqrt{\sigma_{\text{wk}}})}}{4Nz\sqrt{D_{\text{wk}}}} \quad (9)$$

estimated as the time required for a wave with speed Nz (where N is the Brunt-Väisälä frequency and z is altitude) to travel a distance equal to the geometric mean of the cold pool size and the interval between cold pools

270 Finally, the circulation induced in the cold pool parameterizations by the differential heating by convection (and cold pools themselves) contributes as well to the physics tendencies for temperature and humidity. The source terms only contain the vertical convergence of the Reynolds flux associated with mass flux transport (downward in the cold pool and upward in its environment) that reads $\overline{\rho w' \phi'} = 1/g \sigma_{\text{wk}} (1 - \sigma_{\text{wk}}) \delta\omega \delta\phi$ for any conserved variable ϕ so that:

$$\begin{cases} \frac{\theta}{T} Q_1^{\text{wk}} = -C_p \sigma_{\text{wk}} (1 - \sigma_{\text{wk}}) \partial_p (\delta\omega \delta\theta), \\ Q_2^{\text{wk}} = L_v \sigma_{\text{wk}} (1 - \sigma_{\text{wk}}) \partial_p (\delta\omega \delta q_v). \end{cases} \quad (10)$$

Finally, the cold pool ~~models~~-model includes:

- three prognostic variables, derived directly from the model equations: the profiles of $\delta\theta$ and δq_v and σ_{wk} .
- 275 - three diagnostic variables, evaluated from the profile of $\delta\theta$: p_{wk} , C_* and $WAPE$
- three main free parameters: the coefficient k , the density D_{wk} and K_{gw} .

2.4 Triggering and closure of the deep convection scheme

Triggering and closure formulations in LMDZ have been described in Rio et al. (2013). Deep convection is triggered when the Available Lifting Energy (ALE) at cloud base exceeds the convective inhibition (CIN) threshold. This lifting energy is provided either by the thermal plume model (ALE_{th}) or by the cold ~~peels~~pool parameterization (ALE_{wk}):

$$\max(ALE_{th}, ALE_{wk}) > |CIN|. \quad (11)$$

The intensity of the convection depends on the mass flux (M_b) at the cloud base. The closure consists in expressing M_b as a function of an available lifting power ALP , provided by the thermal plume (ALP_{th}) and cold pools (ALP_{wk}) parameterizations:

$$M_b = \frac{ALP_{th} + ALP_{wk}}{(2W_b^2 + |CIN|)}, \quad (12)$$

where

$$W_b = wb_{srf} + \frac{wb_{max}}{1 + \frac{\Delta P}{(P_s - P_{LFC})}} \quad (13)$$

is the vertical velocity at the level of free convection (LFC), $\Delta P = 500$ hPa and wb_{srf} and wb_{max} are model free parameters.

Concerning the boundary layer, the available lifting energy is taken as the maximum kinetic energy in the thermal plume below cloud base

$$ALE_{th} = \frac{1}{2} w_{th,max}^2. \quad (14)$$

A notable improvement was introduced by Rochetin et al. (2014), with the implementation of a statistical representation of the size distribution of cloudy thermal bases. In the new statistical triggering, deep convection is activated if both $ALE_{th} > |CIN|$ and at least one cumulus cloud within a grid cell exceeds a given size, specified by S_{trig} (adjustable parameter). The available lifting power scales with $w_{th,max}^3$.

To calculate ALE_{wk} , the model assumes that the maximum speed (C_{max}) on the cold pool contour will trigger convection. This is assumed to be proportional to the square root of $WAPE$, with a higher coefficient of proportionality than the one used for C_* leading to the following relationship:

$$C_{max} = k' \sqrt{2WAPE}, \quad (15)$$

where $k'=1$.

The Available Lifting Energy associated with cold pools is thus expressed by the following relationship:

$$ALE_{wk} = \frac{1}{2} C_{max}^2. \quad (16)$$

Combining equations (16) and (15), one obtains:

$$ALE_{wk} = k'^2 WAPE, \quad (17)$$

305 with $k' = 1$, meaning that, in the cold pool model, the Available Lifting Energy associated with cold pools is equal to the collapse energy.

Each cold pool provides a power associated with its horizontal spreading. This power is calculated as the product of the kinetic energy supplied by the pool and the mass flux, evaluated over the entire contour of the pool ($L_g = 2\pi r$), over the full height of the pool h_{wk} , and using the spreading velocity C_* . The power ($ALP_{wk,i}$) of an individual cool pool i is therefore:

$$310 \quad ALP_{wk,i} = \frac{1}{2}\rho C_*^3 h_{wk} L_g. \quad (18)$$

To obtain the average power (ALP_{wk}) over all cold pools in the domain, this individual power is multiplied by the cold pool number density D_{wk} :

$$ALP_{wk} = ALP_{wk,i} D_{wk}. \quad (19)$$

Since $\sigma_{wk} = D_{wk}\pi r^2$, the lifting power ALP_{wk} reads:

$$315 \quad ALP_{wk} = \epsilon \rho C_*^3 h_{wk} \sqrt{\sigma_{wk} D_{wk} \pi}, \quad (20)$$

where $\epsilon = 0.25$ is the lifting efficiency. This value means that 25% of the power associated with cold pool spreading is available for deep convection. This rather arbitrary choice is tested hereafter.

2.5 Large Eddy Simulations

Atmospheric Large Eddy Simulations (LES) are performed with non hydrostatic models, with a grid resolution fine enough
 320 to resolve the main structures (large eddies) that dominate the turbulent or convective transport. They provide an explicit and detailed representation of turbulent and convective motions within the boundary layer and of the associated clouds (see e. g. Brown et al., 2002; Siebesma et al., 2003, among many others). LES were used as well for assessing parameterizations of deep convection, including cold pools that are represented explicitly in such simulations (see e. g. Suselj et al., 2019). Note however that these simulations can become more dependent on the model schemes used when microphysics start to play a
 325 predominant role. One of the major strengths of LES lies in its ability to provide three-dimensional information not available from observations, making them an indispensable complement to the latter for understanding processes. In addition, LES can be used to validate the internal variables of parameterizations, enabling their physical realism to be assessed.

In the present study, we use the outputs of LES for two test cases, one over ocean and the other one over land. The test case over ocean was run with two different models, SAM and MesoNH, with the exactly same setup in order to assess the
 330 uncertainty in LES simulations.

The LES over ocean were carried out in Radiative-Convective Equilibrium (RCE) mode (see e. g. Daleu et al., 2015). In the RCE simulations used here, radiative computation is replaced by a constant cooling of -1.5 K per day, while the surface temperature is imposed at 300 K. Two LES of this RCE case are used here, one performed with the SAM model (Khairoutdinov and Randall, 2003) and the other one with MesoNH (Lac et al., 2018). Both simulations cover an horizontal domain
 335 of 200 km×200 km with resolution of 250 m. The lateral boundary conditions are cyclic for both models. These two RCE

simulations ran for 44 days. A quasi steady-state regime is reached after about 40 days. Output are available every 3 hours for SAM and every 24 hours for MesoNH.

The LES over land is issued from AMMA (African Monsoon Multidisciplinary Analysis) field campaign experiment (Re-delsperger et al., 2006). This case is derived from observations made on July 10, 2006 during which a relatively small, short-
340 lived convective system formed over Niamey (Lothon et al., 2011). This system, with a lifetime of around 6 hours, was observed by various instruments (radar and atmospheric soundings), supplemented by satellite data. This case study represents a typical example of deep convection in the Sahel region (Couvreur et al., 2012). An LES of this continental case was carried out with the MesoNH model over a $200 \text{ km} \times 200 \text{ km}$ domain, with a horizontal resolution of 200 m. Lateral boundary conditions are cyclic and surface fluxes are imposed. Outputs are generated at a frequency of 30 minutes.

345 One advantage of RCE for parameterization development is that it targets a steady state regime. By contrasts, the AMMA case corresponds to a diurnal cycle of deep convection over land, for which the time matters, and for which errors in the phasing of the triggering of deep convection can alter comparisons.

3 Assessment of the cold pool model internal equations from LES

3.1 Distinguishing the cold pools from their environment

350 In order to use LES for the assessment of the cold pool parameterization, the first challenge is to separate cold pools from their environment. Indeed, there is no a priori established framework for objectively identifying cold pools in observations and numerical models (Rochetin et al., 2021), and choices may depend in part on the physical picture one has of cold pools, and also, for the purpose at hand, on the picture underlying the parameterization. The first method for identifying cold pools proposed by Young et al. (1995) was based on surface precipitation rates. In more recent studies, such as those by Provod et al.
355 (2016); Zuidema et al. (2017); Vogel et al. (2021); Rochetin et al. (2021); Touzé-Peiffer et al. (2022), the detection of cold pools is closer to a density current oriented detection, in which variations in temperature, pressure and wind are taken into account.

In the present study, the aim is not to isolate individual “cold pools objects”, but only to know whether a grid cell of the LES is inside or outside cold pools. Also the boundary conditions are idealized targeting the statistical homogeneity assumption that
360 is at the basis of the Reynolds decomposition between dynamical core and physics parameterizations. In this idealized case with uniform surface temperature, cold pools can be identified using a threshold on the anomaly (after removing the domain average) of temperature at 10 m above surface, $T_{10\text{m}}$, i.e. at the first model mid layer.

Fig. 2 and 3 show a horizontal moving average with a box of $3.25 \text{ km} \times 3.25 \text{ km}$ of the divergence of the wind at 10 m above surface, $V_{10\text{m}}$. From these maps, the centers and gust fronts of cold pools can be easily identified, corresponding respectively
365 to the maximum and minimum of divergence values. Maxima of divergence of surface wind indicate the center of cold pools where cold air masses collapse. Precipitation is generally co-located with these divergence maxima (not shown). The fairly strong wind convergence observed around cold pools centers corresponds to the strong lift of air masses created upstream of the gust front at the cold pool’s periphery.




Figure 2. Moving average (with a box of $3.25 \text{ km} \times 3.25 \text{ km}$) of the divergence of wind at 10 m (in unit 10^{-3} s^{-1} or 1 m/s/km). Panels a and b correspond to two different states of the LES SAM carried out on the oceanic RCE case. Contours of temperature anomalies at 10 m at -0.4 K (green), -0.2 K (red) and 0 K (black) are superimposed on the smoothed divergence field.

Both the two LES of the RCE case and the LES of the AMMA case show cold pools clusters forming a common gust front.
 370 This can be explained by the fact that, during propagation, cold pools can merge to create a single, larger cold pool. We can also observe that wind convergence (and thus associated updrafts) is more intense where cold pools meet. This is in line with some studies that indicate that convection initiation on gust fronts is more efficient when two or more cold pools collide (Meyer and Haerter, 2020; Torri and Kuang, 2019; Haerter and Schlemmer, 2018; Feng et al., 2015).

We superimpose on this map the $T_{10\text{m}}$ anomaly contours with different values to determine an optimal threshold for this
 375 anomaly. In the RCE case, the $T_{10\text{m}}$ anomaly at 0 K sometimes includes regions without cold pools centers, where divergence of surface wind is low (Fig. 2a and b) while anomaly contours at -0.2 K and -0.4 K surround the centers of cold pools quite well. In the AMMA case, figure 3a clearly shows that the 0 K threshold is too high to identify cold pools. Fig. 3b, on the other hand, shows that the -1 K threshold follows gust fronts of cold pools better than the -0.5 K threshold. On the basis of these analyses, we retain the $T_{10\text{m}}$ anomaly thresholds at -0.2 K and -1 K to identify cold pools in the RCE and AMMA cases
 380 respectively.




Figure 3. Same as Fig. 2 for two successive instants, 5:30 PM (a) and 7:30 PM (b), of the LES MESONH carried out on the AMMA case. The contours superimposed corresponds to T_{10m} anomalies of -1 K (green), -0.5 K (red) and 0 K (black).

3.2 Computing the WAPE from the cold pool anomalies

Once the threshold value is fixed for the T_{10m} anomaly, we separate the full 3-dimensional LES domain between cold pool region (wk) and the rest of the domain (ex) from which we can compute the horizontal averages on each subdomain, X_{wk} inside cold pools and X_{ex} outside, and then the cold pool anomaly $\delta X = X_{wk} - X_{ex}$. This sampling allows to compute the vertical
 385 profiles of cold pools anomaly for temperature (δT), humidity (δq_v) and vertical velocity (δw). Examples of temperature anomalies are shown in Fig. 4.

Note that we apply the same surface mask to the entire column to determine the vertical profiles. This simple vision of vertical cylinders is adopted to match the view underlying the parameterization but may be put into question in the presence of strongly tilted convection.

390 One can then compute the collapse energy (*WAPE*) of cold pools in the LES by integrating from the surface up to p_{wk} , the virtual temperature anomaly, $\delta\theta_v$ (equation, 3). As suggested by Grandpeix et al. (2010), we take for p_{wk} the pressure where the δT profile cancels out. This altitude is around 950 hPa (approximately 600 m) in the oceanic RCE case and around 800 hPa (approximately 2 km) in the AMMA case (Fig. 4).




Figure 4. Vertical profiles of the cold pool temperature anomaly (left, difference between the inside and the outside of cold pools) and of the averaged condensed water (right, g/kg).

3.3 Computing C_* from the mean wind divergence inside cold pools

395 It is assumed in the parameterization that cold pools are identical disks of radius r . This assumption makes it easy to determine C_* by the divergence theorem:

$$\int \int \text{div}(V_{10\text{m}}) dS_{\text{wk}} = C_* L_g. \quad (21)$$

After applying equation 21, we obtain the following relation

$$C_* = \frac{\overline{\text{div}(V_{10\text{m}})} S_{\text{wk}}}{L_g}, \quad (22)$$

400 where S_{wk} is the area of cold pools.

Since $L_g = 2\pi r$, $\sigma_{\text{wk}} = D_{\text{wk}}\pi r^2$ and $S_{\text{wk}} = \pi r^2$, C_* can read as a function of the mean divergence of wind at 10 m, the surface fraction (σ_{wk}) and the density (D_{wk}) of cold pools by the relation:

$$C_* = \frac{1}{2} \overline{\text{div}(V_{10\text{m}})} \sqrt{\frac{\sigma_{\text{wk}}}{D_{\text{wk}}\pi}}. \quad (23)$$

The choice to calculate C_* here from σ_{wk} rather than from the radius r is explained by the fact that it is easier to estimate σ_{wk} in the LES, since we did not use automatic cold-pool detection tools that would allow the radius of cold pools to be determined directly.

To apply this calculation of C_* in the LES, we take the horizontal average of the surface wind divergence inside cold pools. The surface fraction (σ_{wk}) of cold pools calculated in the LES is 0.214 (Average over the available time steps between 5:00 PM and 10:00 PM) for the AMMA case and 0.253 (Average over the 24 time steps with the SAM model) for the RCE case. To determine D_{wk} , we manually counted the centers of cold pools visible on the surface wind divergence maps (Fig. 2 and 3), as we did not use automated detection methods in this study that could generate their number automatically. We find a density, D_{wk} , of about 5 cold pools per $100 \text{ km} \times 100 \text{ km}$ in the RCE case, and about 2.5 cold pools over the same domain in the AMMA case.

3.4 Computing ALP and ALE from gust front vertical velocities

Finally we derive a direct estimation of the Available Lifting Energy (ALE_{wk}) and Power (ALP_{wk}) in the LES from a sampling of the vertical wind at cloud base.

To do this, we first determine an average cloud-base height at which we extract vertical velocities $w_b(x, y)$. This height corresponds to the altitude at which the average profile of condensed water reaches its first non-zero value. It is estimated at around 950 hPa on the two oceanic LES and at around 750 hPa for the LES of the AMMA case (cf. Fig. 4).

We then separate the updrafts on gust fronts from those associated with thermal plumes. Since the updrafts on gust fronts are both stronger and more coherent horizontally than those associated with thermal plumes, we define a gust front mask based on a threshold applied to an horizontally moving average of the vertical velocity at cloud based w_b , denoted as $\tilde{w}_b(x, y)$. Because the gust fronts are stronger in the AMMA case than in the RCE case, different choices were made for the size of the horizontal box of the moving average ($1.25 \text{ km} \times 1.25 \text{ km}$ for the RCE case and $2 \text{ km} \times 2 \text{ km}$ for AMMA) and for the value of the vertical velocity threshold (0.6 m/s for the RCE case and 2 m/s for the AMMA case). Those values were retained after several tests so as to separate as effectively as possible the gust front from other ascents.

Fig. 5 and 6 overlays the updrafts within (red) and outside (green) gust fronts on maps of T_{10m} anomaly (smoothed by applying a moving average with a box of $2.5 \text{ km} \times 2.5 \text{ km}$), for the RCE and AMMA cases respectively. The contours of the T_{10m} anomalies used to identify cold pools (-0.2 K for RCE and -1 K for AMMA) are displayed as well. Visually, the gust fronts computed with $\tilde{w}_b(x, y)$ thresholds of 0.6 m/s (RCE) and 2 m/s (AMMA) align well with the contours of cold pools identified using these T_{10m} anomaly thresholds. It also appears that most thermals are located in the environment of cold pools for both the RCE and AMMA cases. This retrospectively validates a choice made in LMDZ6A, where the effect of thermals was only computed outside cold pools.

Both ALE_{wk} and ALP_{wk} are computed from w_b restricted to the gust front mask, noted $w_{\text{b,gust}}$.

ALE_{wk} is estimated as the kinetic energy associated with the maximum value of $w_{\text{b,gust}}(x, y)$:

$$ALE_{\text{wk}} = \max\left(\frac{1}{2}w_{\text{b,gust}}^2\right). \quad (24)$$

	$D_{\text{wk}}(10^{-10} \text{ m}^{-2})$	σ_{wk}	σ_{gust}
RCE			
LES SAM	5	0.253	0.048
LES MESONH	5	0.264	0.017
AMMA			
LES MESONH	2.5	0.214	0.045

Table 1. Cold pools number density (D_{wk}), surface fraction of cold pools (σ_{wk}), and surface fraction of gust fronts (σ_{gust}) estimated from the LES for the RCE and AMMA cases. For the RCE case, the values represent an average over the 24 available time steps from the SAM LES and the 10 available time steps from the MESONH LES. For the AMMA case, the values are an average of the time steps obtained between 5:00 PM and 10:00 PM.

ALP_{wk} represents the average updrafts power provided by all cold pools in the domain. It is calculated as the horizontal average of the cube of $w_{\text{b,gust}}$ times the surface fraction (σ_{gust}) covered by gust fronts:

$$ALP_{\text{wk}} = \sigma_{\text{gust}} \frac{1}{2} \overline{\rho w_{\text{b,gust}}^3}. \quad (25)$$

440 The gust front mask is used to calculate σ_{gust} , which is 0.048 (LES SAM) for the RCE case and 0.045 for the AMMA case, for the times shown in Fig. 5 and 6. Characteristics of the cold pools estimated from the sampling are gathered on Table 1.

3.5 Validation of phenomenological laws

Physical parameterizations are defined by sets of mathematical equations designed to represent subgrid processes within a column of the model. The formulation of these equations is based both on a phenomenological understanding of the processes
445 involved and on fundamental principles of physics. These parameterizations can be evaluated as a whole or in parts, by isolating certain equations or relationships between internal variables, or between internal variables and GCM state variables. LES offers the possibility of performing a priori validation and adjustment of these laws.

In the cold pool model, variables ALE_{wk} , ALP_{wk} and C_* are determined from the collapse energy, $WAPE$ (see equations (4), (17) and (20)). We compare in Table 2 the values obtained using the parameterization formulations (parameterized value
450 P), based on $WAPE$ deduced from $\delta\theta_v$, with those obtained directly from resolved wind in the LES (sampled value S): the vertical speed at cloud base (w_b) for ALE_{wk} and ALP_{wk} , and the mean divergence of wind at 10 m in cold pools for C_* . These analyses are performed by averaging over the available time steps: 24 time steps for SAM and 7 for MESONH in the RCE case, and between 5:00 PM and 10:00 PM for the AMMA case.

The values of ALE_{wk} calculated by both methods are very close to each other. The largest error is an underestimation by
455 about 30% of the ALE_{wk} computed from $WAPE$ compared to the $w_{\text{b,gust}}$ estimate. These results for the three LES confirm the validity of the hypothesis of equality between ALE_{wk} and $WAPE$ assumed in the parameterization.

Table 2 shows that, C_* values computed from the $WAPE$ are systematically lower than those coming from the mean divergence of wind at 10 m in cold pools. This difference could be due to an underestimation of the coefficient k , imposed




Figure 5. Map of T_{10m} anomaly (color shadings), smoothed (moving average with a horizontal box of $2.5\text{ km}\times 2.5\text{ km}$), at an instant of the LES SAM of the RCE case. The black contour is the -0.2 K anomaly used to separate the inside from the environment of cold pools. The green and red dots show grid cells with vertical velocity at cloud base w_b larger than 0.8 m/s , inside (red) and outside (green) the gust front mask (see main text for the definition of the gust front mask).




Figure 6. Map of T_{10m} anomaly (color shadings), smoothed (moving average with a horizontal box of $2.5 \text{ km} \times 2.5 \text{ km}$), at 7:30 PM for the LES of the AMMA case. The black contour is the -1 K anomaly used to separate the inside from the environment of cold pools. The green and red dots show grid cells with vertical velocity at cloud base w_b larger than 2 m/s , inside (red) and outside (green) the gust front mask (see main text for the definition of the gust front mask).

	ALE_{wk} (J/kg)	$WAPE$ (J/Kg)	C_* (m/s)	C_* (m/s)	C_* (m/s)	ALP_{wk} (W/m^2)	ALP_{wk} (J/kg)	ALP_{wk} (W/m^2)
from:	w_b sampl.	$\int \delta\theta_v$	$\text{div}(V_{10m})$	P: k=0.33	P: k=0.56	w_b sampl.	P: k=0.33	P: k=0.56
RCE SAM	10.460	7.962	2.228	1.315	2.232	0.054	0.008	0.044
RCE MESO	6.965	7.912	2.264	1.313	2.228	0.020	0.008	0.044
AMMA MESO	59.760	45.870	5.362	3.133	5.316	1.733	0.279	1.368

Table 2. Comparison of the variables $WAPE$, ALE_{wk} , C_* , and ALP_{wk} obtained directly from the resolved wind in the LES (from w_b sampling or $\text{div}(V_{10m})$), with those calculated using the formulations of the parameterization (parameterized values P, coming from the $WAPE$ computation based on the $\delta\theta_v$ profile). The S values are derived from the vertical velocity at cloud base (w_b) for ALE_{wk} and ALP_{wk} , from the mean divergence of wind at 10 m in cold pools for C_* , sampled directly from the LES. The P values are calculated from the $WAPE$ deduced from $\delta\theta_v$, itself sampled from the same LES, considering the coefficients $k = 0.33$ and $k = 0.56$. The analyses are based on the average of the available time steps: 24 time steps for the LES performed with SAM and 7 with MESONH in the oceanic RCE case, and between 5:00 PM and 10:00 PM for the LES of the AMMA case.

here at 0.33. With $k = 0.56$, the calculation of C_* based on the $WAPE$ becomes comparable to those obtained from the mean
460 divergence of wind at 10 m in cold pools (Table 2). As discussed above, the value of 0.33 was retained following an oral
communication by Lafore (2000). But other studies propose different values: Lafore and Moncrieff (1989) estimate k at 0.68
based on CRM simulations of 2D squall grain, Bryan (2005) estimate it at 0.5 from observations of cold pools during the
BAMEX experiment in the American Great Plains. Our results are thus compatible with the hypothesis of the model which
postulates that the kinetic energy of cold pools results from the transformation of $WAPE$ into kinetic energy with a coefficient
465 k compatible with the published estimates.

Table 2 also shows that, for the three LES cases, the values of ALP_{wk} calculated with C_* from $WAPE$ are at least three
times lower than those obtained from $w_{b,gust}$. Two coefficients are involved in the calculation of ALP_{wk} with the paramteri-
zation formula: the coefficient k and the lifting efficiency ϵ , imposed respectively to 0.33 and 0.25. Using $k=0.56$ however in
the calculation of C_* , and keeping ϵ at its nominal value of 0.25 allows to reconcile the various estimates. This is compatible
470 with the hypothesis of the parameterization according to which 25% of the horizontal power provided by the cold pools during
its propagation would be used to reinforce the intensity of the convection while a large part dissipates.

4 Evaluation in the single column configuration of LMDZ

In this section, we evaluate the cold pool parameterization in the SCM configuration of LMDZ. The comparison is more
demanding here, since all parameterizations interact with each other and because the state of the atmosphere at the time of
475 evaluation depends on the interaction of all those parameterizations during the preceding hours (AMMA) or days (RCE).

The SCM simulations are performed with exactly the same initial and boundary conditions as the corresponding LES for both cases.

For the RCE case, we represent diagnostics once a quasi-steady state has been reached by averaging results between day 40 and 44.

480 In the AMMA case, cold pools appear around 5:00 PM in the LES but as early as 1:30 PM in LMDZ CTRL, revealing a model limitation. Adjusting the S_{trig} parameter involved in the triggering criteria could delay this onset, though a more physical approach would be required. For comparison, we focus on the times when cold pools are most developed: 7:30 PM in the LES and 2:30 PM in LMDZ CTRL, where the intermediate analysis of δT shows colder and thus more pronounced pools.

In order to facilitate comparisons between LMDZ and LES, we also impose in the LMDZ simulations the density of cold 485 pools estimated in the LES. We thus set a density of 5 cold pools per 100 km \times 100 km for the RCE case and 2.5 cold pools per 100 km \times 100 km for the AMMA case. To represent the profiles of δT , δq_v and δw in LMDZ CTRL for the RCE case, we perform a time average between the 41st and 43rd day of simulation, in order to compare with the LES at the same times. For the AMMA case, the analysis is performed at 7:30 PM in the LES and at 2:30 PM in LMDZ CTRL, as specified above. The same procedure is applied to compare the $WAPE$, ALE , and ALP variables between LMDZ CTRL and the LES for both 490 cases.

4.1 Vertical profiles of δT , δq_v and δw

The analysis of the δT profiles in the LES confirms that cold pools are colder at the surface with temperatures increasing towards the top for the three LES. The cold pools are about three times deeper in AMMA (Fig. 7a) than for the RCE case (Fig. 7d). In the LES, the cold pool temperatures for the AMMA case (around 4 K) are lower than those of the RCE case 495 (around 1.2 K). This is consistent with observations which indicate much colder pools over land than over the ocean. For the AMMA case in particular, observations reveal a temperature drop of approximately 5 K during the passage of the cold pool (Lothon et al., 2011), a value fairly close to that of the LES. It should be noted however that the AMMA case corresponds to a relatively weak episode of continental convection. The δq_v profiles indicate that at the surface, cold pools are wetter than the surrounding air in the RCE case and the AMMA case (Fig. 7b and 7e). In both cases, the excess of humidity within cold pools 500 decreases with altitude up to the cold pools top. The humidity deficit above this level is due to the lateral entrainment of dry air from the mid troposphere and its subsidence into the cold pools (Fig. 7c and 7f). For the RCE case, this subsidence vanishes below 800 hPa (Fig. 7c), while for the AMMA case, it vanishes at a higher level, around 600 hPa (Fig. 7f).

The δT profiles simulated with LMDZ CTRL are qualitatively consistent with LES, with a cold pool top (where δT cancels) at about the right altitude. Cold pools simulated with LMDZ are however warmer than in the LES for the RCE case (Fig. 7a), 505 and colder at the surface than the LES for the AMMA case (Fig. 7d). Consistently with LES, cold pools are also wetter at the surface and drier close to their top top (Fig. 7b and Fig. 7e). However the variations of δq_v are much larger in LMDZ than in the corresponding LES. In particular, the cold pools are much too dry at their top in LMDZ. In both cases, cold pools are associated with subsidence. The height at which the subsidence of air masses into cold pools begins, fixed at 600 hPa in LMDZ CTRL, is too high compared to LES for the RCE case (Fig. 7e).




Figure 7. Vertical profiles of δT , δq_v and δw calculated in the LES and simulated by LMDZ control (LMDZ CTRL). For the RCE case (a, b, c), the profiles are shown for a set of 24 times for the SAM LES (light grey) and 10 for MESONH (dark grey). For LMDZ CTRL, results are averaged from day 41 to 43. For the AMMA case (d, e, f), the profiles correspond to the times when cold pools were most developed, i.e., 7:30 PM in LES and 2:30 PM in LMDZ CTRL.

	$WAPE$ (J/Kg)	ALE_{wk} (J/kg)	C_* (m/s)	ALP_{wk} (W/m^2)
RCE				
LES SAM	7.962	10.460	2.228	0.054
LES MESONH	7.912	6.965	2.264	0.020
LMDZ CTRL	2.957	2.957	0.802	0.001
AMMA				
LES MESONH	62.110	66.960	4.762	2.304
LMDZ CTRL	71.300	75.170	3.941	0.103

Table 3. Comparison of the $WAPE$, ALE_{wk} , C_* and ALP_{wk} computed from sampling of the LES and by LMDZ control (LMDZ CTRL) for the RCE case and the AMMA case. For the RCE case, comparisons are made using an average of the days following the achievement of equilibrium (days 41, 42, and 43). For the AMMA case, they are performed at the times when the cold pools are most developed (7:30 PM in the LES and 2:30 PM in LMDZ CTRL).

510 4.2 WAPE, ALE and ALP

For the RCE case, the $WAPE$ is significantly smaller in LMDZ CTRL than in the LES, with a difference of at least a factor of 2 (Table 3). These low values of $WAPE$ in LMDZ CTRL also translate into low ALE_{wk} values compared to LES (Table 3). On the other hand, for the AMMA case, the $WAPE$ simulated by the model, and consequently ALE_{wk} , are slightly higher than the values derived from the LES (Table 3). The value of C_* simulated by LMDZ CTRL is at least three times smaller than that of the LES in the RCE case and slightly lower for the AMMA case (Table 3). ALP_{wk} is at least twenty times weaker in LMDZ CTRL than in the LES for all cases (Table 3).

Various modifications of the cold pool parameterization are explored in the following section to try to correct the defects listed above.

5 Improvements of the cold pool model

520 Here, we start by correcting the identified discrepancies between the LES and the model concerning the value of the coefficient k and the pressure height p_{upper} , and by assessing the impact of these changes on the temperature and humidity difference between the cold pools and their environment, before exploring other avenues for improvement.

5.1 Coefficient k

We present here the impact of increasing the coefficient k from 0.33 to 0.56 (LMDZ V1 simulation) on the profiles of δT , δq_v , δw as well as on the variables C_* , $WAPE$, ALP_{wk} and ALE_{wk} (see Table 4). In the RCE case, this modification significantly improves the profile of δw below p_{wk} (Fig. 8c). It also allows for a better representation of the δw profiles below p_{wk} in the AMMA case (Fig. 8f). These improvements are directly linked to the increase in C_* for both cases (Table 5), since the




Figure 8. Vertical profiles of δT , δq_v and δw calculated in the LES and simulated with LMDZ: for a control simulation (CTRL, same curves as Fig. 7), with adjustment of the coefficient k to 0.56 (V1), with the modified computation of p_{upper} (V2) and with the activation of thermals in the entire domain (V3). Both the RCE (a, b, c) and AMMA (d, e, f) cases are shown for the times as in Fig. 9.




Figure 9. Vertical profiles of potential temperature (θ) and specific humidity (q_v) calculated in the LES and simulated with LMDZ: for a control simulation (CTRL), with adjustment of the coefficient k to 0.56 (V1), with the modified computation of p_{upper} (V2) and with the activation of thermals in the entire domain (V3). Both the RCE (a, b) and AMMA (c, d) cases are shown for the same times as in Fig. 7

Simulations	Protocols
LMDZ CTRL	simulation of LMDZ with the standard configuration except that D_{wk} is set to 5 per $100 \times 100 \text{ km}^2$ for the RCE case and 2.5 for AMMA
LMDZ V1	LMDZ CTRL + change of k to 0.56
LMDZ V2	LMDZ V1 + changed computation of p_{upper}
LMDZ V3	LMDZ V2 + activation of thermals throughout the domain

Table 4. Description of simulations performed with LMDZ in the standard configuration and with various modifications

δw profile below p_{wk} depends on the spreading of cold pools. The increase in C_* could be associated with stronger air mass subsidence within the cold pool, which would contribute to a slight drying near the surface in both cases (Fig. 8b and 8e). For the AMMA case, this drying results in slightly drier cold pools at the surface in LMDZ V1 than in the LES, but they remain overall comparable to the latter. The increase in C_* in both the RCE and AMMA cases also leads to a better representation of ALP_{wk} (an increase by at least a factor of 5 for both cases), even though this variable remains underestimated (Table 5). We also note a warming effect from this modification in both the RCE and AMMA cases. The impact on the δT profiles in the AMMA and RCE cases is responsible for the decrease in the values of $WAP E$ and $AL E_{\text{wk}}$ for these two cases (Table 5).

Fig. 9 shows that the modification introduced in version V1 has a low impact on the θ and q_v mean profiles (Fig. 9), the black and blue curves being almost superimposed. Both CTRL and V1 simulations reproduce the θ profiles fairly well. Around 600 hPa, the temperature is too warm in the RCE case. Regarding humidity, a dry bias is present in the boundary layer in the RCE case, as well as between 800 and 400 hPa. For the AMMA case, there is a wet bias in the boundary layer and above 600 hPa, and a dry bias between 700 and 600 hPa. At the altitudes where these dry biases are found, notably between 800 and 400 hPa for the RCE case and between 700 and 600 hPa for the AMMA case, LMDZ produces almost no clouds, likely due to an overly dry atmosphere (not shown).

5.2 Choice of cold pools scheme upper bound, p_{upper}

In the previous sections, we found that the altitude at which the subsidence of dry air above cold pools initiate is located around 800 hPa in the LES for the RCE case and around 600 hPa for the AMMA case, while in LMDZ, p_{upper} is arbitrarily set to 600 hPa in the original version of the parameterization. In version V2, in addition to the change of the value of k from 0.33 to 0.56, we impose $p_{\text{srf}} - p_{\text{upper}} = \gamma_{\text{wk,upper}}(p_{\text{srf}} - p_{\text{wk}})$ with $\gamma_{\text{wk,upper}}$, fixed here to 3, is considered as a new free parameter in the following section.

Note that the numerical scheme used to estimate p_{wk} was also modified compared to GL10 in order to solve instability issues.

The principle of the new scheme is to compute the height at which the vertical integral of $\delta\theta$, from the surface to that level, reaches a fraction χ of the total integral from the surface to the first positive value of $\delta\theta$. This fraction is taken slightly below 100% to avoid fluctuations of p_{wk} from one time step to the other, which happens when $\delta\theta$ is close to zero on a large range of

altitudes around p_{wk} as seen in Fig. 7 and Fig. 8. The new computation was activated already in CTRL and V1, not affecting the results significantly (figure not shown). The new computation becomes really important when p_{upper} is deduced from p_{wk} rather than being imposed.

Comparisons between LMDZ V2 simulations and LES show a better representation of the δq_v profiles at the top of cold pools in both the RCE and AMMA cases (Fig. 8b and 8e). These results show that the dry bias at the top of the cold pool in the original version was due to advection of dry air from too high an altitude. This modification also reduces slightly the humidity of cold pools near surface in the RCE case, although they remain more humid than in the LES. Note finally that this modification has a very limited impact on the δT profiles.

Table 5 shows that the change in p_{upper} weakly affects the variables $WAPE$, C_* , ALE_{wk} and ALP_{wk} for these two cases. Versions V2 does not modify much the mean vertical profiles except for a drying of the mid-troposphere in the RCE case (Fig. 9b), in a region where the CTRL simulation was already too dry. Although the time evolution of the mean profiles is the first target of physics parameterizations, we think however that the improvement of the internal variables is so strong for this modification that it should be adopted in the future in LMDZ.

5.3 Activation of thermals throughout the domain

As explained above, in the standard LMDZ configuration, thermals only interact with temperature and humidity profiles outside cold pools, inducing a differential heating in moistening (equation (6)). This choice was originally made to account for the fact that the atmosphere is more stable inside cold pools, and indeed the analysis above shows that the thermal plumes that reach cloud base are essentially located outside cold pools. Version V3 is identical to version V2, except that we consider that thermal plumes are active everywhere in the grid cell. Consistently, the terms Q_1^{th} and Q_2^{th} are removed from (6). For the RCE case, this leads to a decrease in the surface humidity of cold pools, closer to the LES results (Fig. 8b). In the AMMA case, the effect is also present, although less pronounced (Fig. 8e). This result is expected because the vertical transport by thermals systematically dries the surface (Diallo et al., 2017). This confirms the key role of boundary layer convection in regulating surface humidity on both continent (Diallo et al., 2017) and ocean (Hourdin et al., 2020), via the mixing of moist air with dry air above. One way to improve the representation of humidity anomaly profile without activating the thermal plumes everywhere would be to add a simple parameterization of shallow and cloud-free boundary layer convection (a simplified version of the thermal plume model) within the cold pool region.

In version V3, cold pools are colder than in version V2 in both the RCE and AMMA cases (Fig. 8). In the RCE case, however, cold pools remain less cold than in the LES despite this effect. In the AMMA case, this cooling accentuates the overestimation of the cold anomaly. In both cases, this cooling leads to an increase in the $WAPE$, C_* , ALE_{wk} , and ALP_{wk} variables (Table 5). Version V3 does not introduce any notable changes, compared to V2, in the profiles of θ and q_v for the RCE and AMMA cases.




Figure 10. Like Fig. 8, but showing for LMDZ, the V2 simulation, the 12 best simulations from the tuning (TUNE, in black) as well as the best one among them (TUNE BEST, in violet dotted).




Figure 11. Like Fig. 9, but showing for LMDZ, the V2 simulation, the 12 best simulations from the tuning (TUNE, in black) as well as the best one among them (TUNE BEST, in violet dotted).

	$WAPE$ (J/Kg)	ALE_{wk} (J/kg)	C_* (m/s)	ALP_{wk} (W/m^2)
RCE				
LES SAM	7.962	10.460	2.228	0.054
LES MESONH	7.912	6.965	2.264	0.020
LMDZ CTRL	2.957	2.957	0.802	0.001
LMDZ V1	2.663	2.663	1.292	0.004
LMDZ V2	2.620	2.620	1.282	0.004
LMDZ V3	3.585	3.585	1.500	0.0055
12 bests	[5.1,5.3]	[5.1,5.3]	[1.79,1.83]	[0.013,0.025]
AMMA				
LES MESONH	62.110	66.960	4.762	2.304
LMDZ CTRL	71.300	75.170	3.941	0.103
LMDZ V1	56.930	56.930	5.975	0.234
LMDZ V2	57.990	57.990	6.031	0.252
LMDZ V3	58.190	58.190 <td 6.041	0.334	
12 bests	[40,60]	[40,60]	[4.5,5.1]	[0.5,2.5]

Table 5. Comparison of the variables $WAPE$, ALE_{wk} , C_* and ALP_{wk} calculated from the samplings in the LES, with those obtained with LMDZ: in a control simulation (CTRL), with the adjustment of the coefficient k to 0.56 (V1), with the modified computation of p_{upper} (V2), the activation of thermals in the entire domain (V3) and for the 12 best simulations of a tuning experiment, for the RCE and AMMA cases.




Figure 12. Time series of the collapse energy, $WAPE$ (J/kg, panel a), and the available lifting power, ALP_{wk} (W/m^2 , panel b), associated with cold pools, shown for the 12 best simulations from the tuning (TUNE, in black), for the best among them (TUNE BEST, in violet dotted) and for the LES (in grey) over times steps between 5:00 PM and 10:00 PM

5.4 Tuning of free parameters

585 The tests presented above show possible avenues for improving the cold pool parameterization. However, we see that the modifications do not sufficiently affect the mean profiles to reduce these biases significantly. All tests underestimate (for the RCE case) or overestimate (for the AMMA case) the cold temperature anomaly inside cold pools, as well as $WAPE$, ALP_{wk} , C_* , and ALP_{wk} . We also observe systematic errors in the mean profiles, notably profiles that are much too dry for the RCE case.

590 In the GCM, these variables are not sensitive only to the parameters or formulation of the cold pool model. They are influenced by all other parameterizations, and in particular by the convection scheme to which the cold pool scheme is strongly coupled. In an attempt to see how modifications to other parameterizations could help reduce these biases, we performed automatic calibration simulations using the `htexplo` tool.

In practice, we decided to start from a tuning performed for the convective boundary layer by Hourdin et al. (2021), for
595 a configuration with 95 levels rather than 79 and using more recent versions of the codes than those used in the rest of the paper. This version indeed serves as the basis for preparing the future version of the climate model for the FastTrac part of the CMIP7 project, whose simulations are scheduled to begin in early 2026. We assume that the boundary layer parameters [values](#) have already been ~~optimized~~ [chosen](#) to accurately represent the convective boundary layer and associated clouds, cumulus and stratocumulus. Regarding modifications to the cold pool model, those affecting the coefficient k and p_{upper} are taken into
600 account, as in the V2 configuration. Adjustments related to thermals (V3) are not considered here because they raise as many questions as they solve.

The tuning is performed using the `htexplo` tool as explained in Appendix A. We choose metrics preferentially on the RCE case targeting the WAPE, ALP and the profiles of anomalies and mean variables. Indeed, we wish to avoid being overly dependent on errors in the phasing of the deep convective diurnal cycle. The adjustable parameters selected for the cold pool
605 model are: k , $\gamma_{wk,upper}$, χ , and ϵ . Parameters involved in the deep convection scheme are also included: the minimum wb_{srf} and maximum wb_{max} vertical velocities at the base of the convective column; the fraction of the grid cell area in which precipitating downdrafts occur, σ_{desc} ; and the maximum precipitation efficiency in the Emmanuel scheme (EP_{max}). This efficiency is a maximum efficiency at the top of the convective columns. The difference $1-EP_{max}$ controls how much condensed water exits the convective clouds, and thus the moisture source in the upper atmosphere. Details on the metrics and parameter ranges
610 chosen are given in Appendix A.

The result of this tuning is the product of ~~extensive~~ trial and error [preliminary tuning experiments](#) regarding the choice of parameters, their bounds, the metrics to adjust, and the associated tolerances. We [finally perform 13 waves of tuning with the experimental setup described in Appendix A, and](#) present here the 12 best ~~simulations resulting from configurations among all the runs performed during this tuning experiment. The best simulations are selected according to the 13 waves of tuning~~
615 ~~Among them, the simulation considered to be the most performant (TUNE-BEST) is also identified.~~ [metrics involved in the tuning procedure \(cf. Table A1\).](#) For each metric we compute a score $s_i = (m_i - o_i) / \sigma_i$ where m_i is the metrics computed on

the simulation, o_i the target value ("observation") and σ_i a tolerance to error. A combined metric is computed as the maximum of s_i over the 13 metrics: $S_i = \max_i s_i$. The best simulations are those with the smallest values of S_i .

The analysis of the results for the RCE case reveals that the ~~simulations improve the representation of 12 bests configurations~~
620 better simulate the targeted variables, particularly the mean humidity profile and the amplitude of the potential temperature deviations $\delta\theta$. These deviations are more negative, consistent with a stronger WAPE and C_* coefficient. Furthermore, the δq_v profiles at the top of the cold pools, as well as the δw profiles, remain well represented across all 12 simulations.

By applying the parameters from the tuning performed on the RCE case to the AMMA case simulations, a significant reduction in the cold bias at the surface of cold pools is obtained (Fig. 10d), as well as an improvement in the mean humidity
625 profiles across all 12 simulations. Only the ~~results from the BEST TUNE simulation reproduce best simulation (BEST TUNE)~~
reproduces δq_v and δw profiles consistent with the LES, the others tending to generate cold pools that are too dry at their top due to slightly overestimated δw profiles (Fig. 10e and 10f). The BEST TUNE simulation also provides a better representation of WAPE (Fig. 12a) and ALP_{wk} (Fig. 12b) in the AMMA case, with values very close to those from the LES at the time when cold pools reach their maximum development (Table 5), despite a shift by about 3 hours earlier in the afternoon. ~~This~~
630 ~~premature triggering in the LMDZ model for the AMMA case has already been showed in Rio et al. (2009) and represents a known limitation of the model. A parameterization of-~~

It is well known that representing the diurnal cycle of convection remains a major challenge for GCMs. In LMDZ, a significant improvement in the representation of the diurnal cycle of precipitation was achieved thanks to the preconditioning of convection by thermals, as well as the role of cold pools in sustaining convection (Rio et al., 2009). Nevertheless, the delay
635 in the transition between shallow and deep convection, currently prescribed according to a cumulus size threshold as discussed above, could correct this bias. Nevertheless, despite this shift in convection triggering, the BEST TUNE simulation shows good agreement with the LES in terms of the duration and intensity of convection. Nevertheless, a moist bias and a cold bias persist in the boundary layer. timing of convective initiation remains a limitation. Intermediate tests have shown that tuning the parameter S_{trig} , which defines the threshold cumulus size required to trigger deep convection, can delay convective onset.
640 A work aiming at better representing the transition between shallow and deep convection is currently ongoing in the LMDZ team.

Several lessons can be learned about the range of the free parameters involved in the tuning when considering the best values (last column of Table B1).

First the fraction of the grid cell assigned to unsaturated downdrafts, σ_{desc} with a selected range of [0.042, 0.048], is larger
645 than the nominal value of 0.015, which contributes to first order to the increased intensity of cold pools, in particular for the RCE case. Concerning the parameters of the cold pool model itself, first the tuning of k from (4) confirms the findings of Section 3.5, with selected range of [0.56, 0.57]. The value of $\gamma_{wk,upper}$ that controls the height of the cold pool model is [3.6, 4.1], compatible as well with the previous findings. With a range of [0.26, 0.46], the tuning suggest that more than 25% of the power associated with cold pool spreading is available for deep convection. Note finally that the χ parameter of the
650 new numerical scheme for p_{wk} computation is also strongly constraint by the tuning exercise (to a value of about 0.97). ~~This exercise confirms the extraordinary progress represented by the availability of automatic retuning methodology.~~

655 It would be difficult to attribute this results in BEST-TUNE simulation to a specific parameter. The calibration result actually arises from a combination of the values of the different parameters used for tuning. Even if a parameter has been significantly modified, its effect on the final result still depends on the other parameters. However, during intermediate tests, we found that the cold pool model is highly sensitive to the parameter σ_{desc} . An increase in its value. The `htexplo` tool retrieves automatically cold pool model parameter values close to those issued from direct analysis of LES. This highlights both the power of using such a tool for model development, avoiding a long phase of trial error in a multi-dimensional space, and the relevance of the model physics. Likewise, the range of values selected for σ_{int} leads to stronger cooling of the cold pool heights of about 600 m in the RCE case and 2 km in the AMMA case, in agreement with the expected orders of magnitude over ocean and over land, respectively.

660 One goal of the tuning is to intensify cold pools, particularly in the RCE case. The fact the increase of σ_{desc} is primarily responsible for strengthening cold pools in this case through stronger evaporation increases confidence in the validity of the physics implemented in the cold pool model. In the AMMA case, the opposite effect (less colder cold pools) is explained by the rather large increase in C_* (from 3 to 5 m/s), induced by the adjustment of k . This increase in C_* accelerates the increase of the cold pools fractional area σ_{wk} , reducing the efficiency of evaporative cooling, which can no longer compensate for the dilution due to their rapid evolution. Although this mechanism is also present in the RCE case (with an increase in C_* from about 1 to 2 m/s), it remains much less pronounced there, allowing the evaporation obtained with these σ_{desc} values to effectively strengthen the cold pools.

670 We also consider that the increase of $1-EP_{max}$ and σ_{desc} played an important role in the improvement of the specific humidity profiles for the RCE and AMMA cases, although other parameters may also play a role. The increase of the difference $1-EP_{max}$ increases the amount of condensed water released at the top of the convective columns, thereby moistening the upper layers of the atmosphere. The value of EP_{max} was set to 0.999 in the control configuration. The increase in σ_{desc} may both enhance humidity through enhanced evaporation and dry the boundary layer, for the AMMA case in particular, due to increased entrainment and downward advection of dry air from the free troposphere.

675 We also briefly examined the behavior of updrafts, given their strong coupling with downdrafts and cold pools. We therefore assume that this parameter played a key role in the BEST-TUNE results note a decrease in undiluted adiabatic updrafts (which represent the theoretical capacity of the column to transport air and moisture) and a decrease in saturated updrafts in the tuned simulations (not shown). These effects are more pronounced in the AMMA case. The decrease in undiluted adiabatic updrafts would be linked to the drying of the boundary layer, itself associated with the increase in σ_{desc} , as explained previously. This reduction in undiluted updrafts would also contribute to the decrease in saturated updrafts by limiting the moisture supply to the mixing with environmental air. In addition, the increase **increase ou decrease ??? Est-ce que c'est ferme tout ça ? Ce paragraphe pourrait peut être n'être que dans la réponse aux reviewers** in the export of condensed water out of convective clouds, which enhances the drying of the column, could also lead to a lower degree of saturation within the convective column.

685 Je pense qu'on peut enlever cette phrase ici car elle est très bien reprise dans la conclusion : All of these results highlight the power of the `htexplo` tool, not only for retuning parameters but also for understanding the role of the different parameters and their interactions.

OK pour affirmer ça mais il faudrait être plus explicite sur les changement de paramètres entre la simulation de contrôle et les valeurs retenues par le tuning. Les donner pour tous les paramètres dont tu parles. Par exemple donner : la valeur de la simulation de contrôle, la plage pour les 12 et la valeur de la best

690 6 Conclusions

This study presents ~~, for the first time, a detailed an~~ evaluation of the ~~cold-pool model Grandpeix and Lafore (2010) within the LMDZ climate model, LMDZ cold pool model~~ based on Large Eddy Simulations (LES). We evaluate both the fundamental assumptions of the model, ~~its internal variables and those involved in the~~ the properties of the cold pools it simulates, as well as its coupling with deep convection, ~~based on two oceanic LES in the RCE regime and a continental LES of using LES of~~ two test cases: the RCE case offers a controlled and quasi-stationary framework for studying convective processes, whereas the AMMA case ~~. For this~~ corresponds to a typical case of Sahelian deep convection where the time dependence matters.

For evaluation, we introduce two ~~samplings kinds of sampling~~ of the LES. The first one separates the interior from the outside of the cold pools, based on a threshold of the 10 m temperature anomaly of -0.2 K for the RCE case and -1 K for AMMA. The second one separates the zone of gust fronts by smoothing horizontally the vertical wind at cloud base (moving average over a square of 1.25 km side for the RCE case and 2 km for AMMA) with a threshold of 0.8 m/s for the RCE case and 2 m/s for AMMA. The coincidence of the temperature contour used for the cold pool sampling with the lines of maximum wind converge near the surface and with the gust front mask provides a very consistent view of the cold pools. It reinforces the choices that guided the conception of the cold pool scheme. It also confirms that most of the thermals reaching cloud base are located outside cold pools. ~~We started by validating relationship~~

We first evaluated directly in the LES relationships internal to the ~~parameterizations, by diagnostics of the LES. This validation mainly relies on comparisons between LES. The cold-pool model variables, diagnosed within the LES according to the formulation of the parameterization, were compared with those obtained directly through sampling in the same simulations~~ parameterization by comparing the value of cold pool spread mean horizontal velocity C_s , obtained from the mean divergence of the horizontal wind within the cold pools, the $WAPE$ computed from the air density contrast between the cold pool and its environment and the lifting energy (ALE) and power (ALP) obtained from the vertical velocity at cloud base. The good consistency between the results confirms the fundamental assumptions of the model. ~~The discrepancies observed for some variables are mainly explained by the choice of a free parameter value, which we adjusted in agreement with the estimates provided by the LES. The validation~~

The evaluation was then carried out in the SCM version of the LMDZ model for the AMMA and RCE cases. In this framework, validation evaluation is much more demanding due to the influence of other parameterizations. ~~The LMDZ simulations were performed with initial and boundary conditions strictly identical to those of the LES.~~

Overall, the comparisons ~~demonstrate the relevance confirm the ability~~ of the model to reproduce the properties of cold pools, although some discrepancies ~~remain. The sources of these differences have been identified, and improvements have been implemented. The dry bias simulated at the top of cold pools is~~ with respect to the LES remain. Part of the discrepancies

720 ~~with LES were attributed to the pressure-of-the-maximum-subsidence-height choice of free parameters rather than to the physical~~
~~formulation of the cold pool model itself. In addition, the computation of the altitude (p_{upper}) which was imposed at a fixed~~
~~value-of-at which the subsidence of dry air within the cold pool vanishes was significantly modified. Initially set to 600 hPa in~~
~~the original scheme. This value which was inspired from continental situations is well suited for the AMMA case but much to~~
~~small-over both ocean and land, this value proved to be too high for the RCE case-for which the LES indicate a value-of-about~~
725 ~~800 hPa. By making~~, leading to excessively dry conditions near the top of the cold pools. From now on, this altitude p_{upper}
~~dependent-on~~ p_{wk} , the pressure-at is determined as a function of the cold pool top , we significantly improve the simulated
~~humidity at the top of cold pools for both cases. These results highlight the control of the cold pool humidity anomaly profile~~
~~by height (p_{wk}), allowing its regional variability (ocean or land) to be taken into account. The importance of this determination~~
~~of p_{upper} highlights the importance of the representation of the~~ lateral entrainment of ~~dry air from the mid-troposphere and~~
730 ~~subsidence-within-free tropospheric dry air into~~ the cold pool .They also confirm the relevance of the cold-pool-transport-model
~~with lateral entrainment between p_{upper} and p_{wk} feeding the maximum subsidence at p_{wk} . A wet bias is also obtained at the~~
~~surface of cold pools in the RCE and AMMA cases. Our analyses show that this bias is linked to the absence, in the model,~~
~~of the effect of thermals on the variation of humidity at the surface of cold pools. The evaporation flux plays a weak role in~~
~~this variation, which seems to be mainly controlled by thermals. Warm and cold biases were also observed at the surface of~~
735 ~~the cold pools, respectively for the RCE and AMMA cases. These biases led to a poor estimation of certain cold-pool-model~~
~~variables ($WAPE$ and C_*), as well as those involved in the coupling, notably ALP_{wk} , since these variables strongly depend~~
~~on the temperature of the cold pools in the model. In order to check whether this limitation may come from a coupling with~~
~~the other model parameterizations, and in particular that of deep convection, we conducted a calibration experiment using the~~
~~HighTune explorer software to jointly adjust the free parameters of the cold-pools and deep convection models. This tuning~~
740 ~~procedure also aimed to correct the dry and wet biases still present in~~ for controlling in particular its humidity.

Thanks to the adjustments made with the potential temperature and specific humidity profiles. These adjustments led to a
~~significant improvement in the representation of cold pool temperature, as well as specific humidity for the RCE and AMMA~~
~~cases, even if a humid and cold bias persists in the boundary layer for~~ htexplo tool, taking into account certain parameters
~~of deep convection, the representation of cold pool temperature has been improved. Analyses indicate that this improvement~~
745 ~~is mainly due to the surface fraction parameter σ_{desc} associated with precipitating downdrafts. Indeed, an increase in this~~
~~parameter leads to more intense evaporation, which enhances the cooling of cold pools. The calibration experiments also~~
~~significantly improved specific humidity in the AMMA and RCE cases. On the one hand, the increase in σ_{desc} corrects the~~
~~moist bias in the boundary layer by drying it out through an increased supply of dry air from upper levels. On the other~~
~~hand, the more intense evaporation linked to the increase in σ_{desc} together with the increase in the difference $1 - EP_{max}$~~
750 ~~moisten the atmosphere above the boundary layer, thereby reducing the dry bias that was present there. Indeed, the increase~~
~~in the difference $1 - EP_{max}$ corresponds to a greater release of condensed water into the upper layers of the atmosphere.~~
~~These results highlight the ability of the htexplo tool to propose a physically consistent calibration while improving the~~
~~understanding of the role of parameters and their interactions.~~

For the AMMA case.~~The values of $WAPE$, ALP_{wk} and C_* are also improved.~~

755 ~~The model also captures~~, the analysis of the diurnal cycle also shows that the adjusted model reproduces well the intensity and duration of convection in the AMMA case, although updraft power of cold pools at the time of their maximum development, but with a delay of about ~~three hours is observed in the onset~~ 3 hours in the diurnal timing of this maximum. This delay may be due to a physical limitation in the transition from shallow to deep convection. Since the in convective initiation is a known limitation of LMDZ, even though the representation of the diurnal cycle of continental precipitation has significantly improved thanks to the inclusion of cold pool effects (Rio et al., 2009). The remaining delay in convective onset can not be modified by the cold pool model itself, since the cold pool model is by essence not active before convective initiation. It is most probably a question of transition between shallow and deep convection and specific work are ongoing in the LMDZ team to better represent this transition.

760 The LES of the AMMA case ~~corresponds to relatively organized deep convection, it allowed also for analyzing the model behavior in this type of regime. In such situations, cold pools strongly interact with wind shear and mesoscale organization, which may alter their dynamics. This is particularly important, given that a parameterization of deep convection organization is currently under development in LMDZ, even though using LES of~~ is characterized by a significant wind shear resulting in relatively organized convection. Although the interaction between wind shear and cold pools is not yet explicitly taken into account in LMDZ, the results indicate that the cold pool model already represents quite well regimes with moderate wind shear. Accounting for wind shear in the parameterization of both convection and cold pools is under consideration in the LMDZ team. Once such effects taken into account, it may be relevant to recalibrate the cold pool model using, in addition to the RCE and AMMA cases, LES of strongly organized convection such as squall lines or ~~mesoscale convective systems~~ Mesoscale Convective Systems (MCS), ~~characterized by highly organized convection, could provide stronger constraints on the cold pool model~~, for which the influence of wind shear is more pronounced.

775 Calibration performed in SCM is not a general guarantee of improvement in 3D mode. It could even degrade the results if the cases used in SCM mode are not enough representative, or if coupling with the large scale (not accounted for in SCM) matters more than the details of the scheme, which could motivate tuning in intermediate configurations before full 3D implementation (Ma et al., 2015). In the present case, however, preliminary 3D tests (not shown) run the most recent versions of the LMDZ climate model show a behavior that is overall similar to that obtained in SCM. The new numerical scheme proposed for the computation of p_{wk} also proved to be more robust, significantly reducing crashes during 3D simulations. The above mentioned changes have been adopted in the new version of the LMDZ global model, used as the atmospheric component of the IPSL-CM7 coupled model under development for the forthcoming CMIP7 Fast-Trac exercise.

785 Although significant progress has been made in recent years in modeling cold pools, due to their important role in convection, challenges remain. First~~a~~, as already mentioned, a missing physical process has been identified in the cold pool model: not accounting for boundary layer convection within the cold pools results in overly humid conditions at the surface. A simple parameterization of boundary layer convective transport, based for instance on a simplified version of the thermal plume model, could be included to better represent vertical mixing within the cold pools without activating the thermal plume model uniformly over the grid cell. The cold pool number density should become an internal variable of the model since we know it presents very different values when considering popcorn like convection over ocean or continents, or well organized long

790 live system such as squall lines. A parameterization of this number density, based on a population dynamic model is presently under test. To end with, the issue of the propagation of cold pools from grid cell to grid cell needs **to be also integrated into GCMs**should also be added to the model.

Code and data availability. The codes used are distributed openly through `subversion`. The LMDZ GCM is available at:

`http://svn.lmd.jussieu.fr/LMDZ/LMDZ6/trunk`

795 It can be installed automatically on a laptop with the command `http://lmdz.lmd.jussieu.fr/pub/install.lmdz.sh` The `htexplo` tool is available at:

`https://svn.lmd.jussieu.fr/HighTune/trunk`

The data and scripts used to make the figures of the document will be made available as well on a DOI if the article is accepted for publication.

The scripts that will be made available include downloading the correct versions of the LMDZ and `htexplo` software.

800 **Appendix A: and the tuning setup**

A1 High-Tune Explorer (`htexplo`) tool

The tuning experiments shown here are done with the `htexplo` tool.

`htexplo` has been developed in collaboration between the LMD (Paris), the Centre National de Recherche Météorologiques (CNRM/Météo-France) and the University of Exeter (UK). It is an automatic calibration tool for free parameters, based on
805 machine learning techniques from the uncertainty quantification community (Williamson et al., 2013). This approach proposes a new calibration paradigm: instead of optimizing parameter values, it aims to identify the subset of parameters that enables the model to reproduce certain observables to a certain accuracy. The main steps involved in using the tool, as well as its mathematical foundations, are well described in Couvreur et al. (2021). The `htexplo` tool was used for the first time in a SCM/LES comparison on several boundary layer cases of the LMDZ model, in order to characterize the subspace of free
810 parameter values for which SCM simulations are consistent with LES for certain metrics and a given tolerance (Couvreur et al., 2021). This information was then used by Hourdin et al. (2021) to calibrate the 3D configuration. These authors demonstrated how reducing the parameter space using this method significantly saves computing and human resources. They also pointed out that this approach eases the burden on the modeler, enabling him or her to concentrate more on understanding and improving the physical parameterizations of the model.

815 For the RCE case, we target the quasi-equilibrium phase by considering averages between days 41 and 43. The metrics selected for these calibration exercises are the profiles of δT , q_v , and θ , evaluated from vertical averages at different levels as indicated in.

Details on the metrics, with targets and tolerances to error, are given in Table A1.

The parameters chosen for tuning are listed in Table B1 together with the a priori ranges given to `htexplo` at the beginning
820 of the tuning exercise and ranges of the best 12 simulations obtained after 13 waves of tuning.

metric	unit	target	tolerance
RCE case, average from day 41 to day 43			
WAPE	$\text{m}^2 \text{s}^{-2}$	8	2
C_*	m s^{-1}	2.2	0.2
$\delta\theta_{0-50 \text{ m}}$	K	-0.83	0.045
$\delta\theta_{0-600 \text{ m}}$	K	-0.48	0.063
$q_v_{0-500 \text{ m}}$	g/kg	14.1	0.45
$q_v_{1-3 \text{ km}}$	g/kg	9.14	0.45
$q_v_{5-6 \text{ km}}$	g/kg	2.55	0.33
$q_v_{8-10 \text{ km}}$	g/kg	0.289	0.063
$\theta_{0-500 \text{ m}}$	K	296.5	1.47
$\theta_{1-3 \text{ km}}$	K	301.0	1.
$\theta_{5-6 \text{ km}}$	K	317.4	1.
$\theta_{8-10 \text{ km}}$	K	328.8	3.
AMMA case, average from 10:00 AM to 5:00 PM			
WAPE	$\text{m}^2 \text{s}^{-2}$	20	3

Table A1. Metrics (targets and $1-\sigma$ tolerances) used for the tuning. For the RCE case, they concern the averages between days 41 and 43 of the WAPE and the cold pool spreading rate C_* , as well as the vertical profiles of $\delta\theta$, q_v , and θ averaged over the altitude ranges specified in the right column. For the AMMA case, only the WAPE averaged between hours 10 and 17 of the simulation is used.

Author contributions. Mamadou Lamine Thiam: work design, diagnostics, analysis, figures and writing

Frédéric Hourdin: work design, diagnostics, analysis and writing

Jean-Yves Grandpeix: conception of the work, analysis and writing

Catherine Rio: analysis and writing

825 Maëlle Coulon–Decorzens: tuning expertise

Competing interests. None

Disclaimer. None

Parameter	units	[min,max] prior	exploration	[min , max] 12 best simulations
Cold pool model				
ϵ ((20))	-	[0.25 , 0.5]	linear	[0.26 , 0.46]
$\gamma_{wk,upper}$	-	[3.5 , 5]	linear	[3.6 , 4.1]
k ((4))	-	[0.33 , 0.66]	linear	[0.56 , 0.57]
χ	-	[0.75 , 0.99]	linear	[0.96 , 0.987]
Convection model				
wb_{srf}	m/s	[0.5 , 1.2]	linear	[0.55 , 0.98]
wb_{max}	m/s	[2.8 , 6]	linear	[2.8 , 3.5]
σ_{desc}	-	[0.015 , 0.05]	linear	[0.042 , 0.048]
$1-EP_{max}$	-	[0.05 , 0.1]	log	[.93 , .95]
$k_{ALP,BL}$	-	[0.2 , 0.5]	linear	[0.33 , 0.47]

Table B1. Free parameters considered in the tuning exercise.

Acknowledgements. The authors thank Caroline Müller for providing the RCE simulation run with the SAM model and Fleur Couvreur for providing the AMMA simulation run with the MESO-NH model. We also acknowledge support from the DEPHY research group of CLIMERI, funded by CNRS/INSU and Météo-France. The work is part of and was supported by the DEPHY french national project. It benefited from the development of the `htexplo` tool within this project and automation of LES/SCM simulations through standardization of IO formats. Mamadou Lamine Thiam’s doctoral thesis was supported by Institute de Recherche pour le Développement (IRD) as part of the international doctoral program entitled “Modélisation des Systèmes Complexes” (PDI-MSc). The thesis also received financial support from the Cuomo Foundation (Monaco) as part of an IPCC support program. [This study has received funding from Agence Nationale de la Recherche - France 2030 as part of the PEPR TRACCS programme under grand number ANR-22-EXTR-0008.](#)

References

- Böing, S. J., Jonker, H. J., Siebesma, A. P., and Grabowski, W. W.: Influence of the subcloud layer on the development of a deep convective ensemble, *Journal of the Atmospheric Sciences*, 69, 2682–2698, 2012.
- Boucher, O., Denvil, S., Levavasseur, G., Cozic, A., Caubel, A., Foujols, M.-A., Meurdesoif, Y., Balkanski, Y., Checa-Garcia, R., Hauglustaine, D., et al.: IPSL IPSL-CM6A-LR-INCA model output prepared for CMIP6 RFMIP piClim-aer, 2020.
- 840 Brown, A., Cederwall, R. T., Chlond, A., Duykerke, P. G., Golaz, J.-C., Khairoutdinov, M., Lewellen, D., Lock, A., MacVean, M., Moeng, C.-H., et al.: Large-eddy simulation of the diurnal cycle of shallow cumulus convection over land, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 128, 1075–1093, 2002.
- Bryan, G. H.: Spurious convective organization in simulated squall lines owing to moist absolutely unstable layers, *Monthly weather review*, 845 133, 1978–1997, 2005.
- Charba, J.: Application of gravity current model to analysis of squall-line gust front, *Monthly Weather Review*, 102, 140–156, 1974.
- Chatfield, R. B. and Brost, R. A.: A two-stream model of the vertical transport of trace species in the convective boundary layer, *J. Geophys. Res.*, 92, 13 263–13 276, 1987.
- Couvreux, F., Rio, C., Guichard, F., Lathon, M., Canut, G., Bouniol, D., and Gounou, A.: Initiation of daytime local convection in a semi-arid 850 region analysed with high-resolution simulations and AMMA observations, *Quarterly Journal of the Royal Meteorological Society*, 138, 56–71, 2012.
- Couvreux, F., Hourdin, F., Williamson, D., Roehrig, R., Volodina, V., Villefranche, N., Rio, C., Audouin, O., Salter, J., Bazile, E., et al.: Process-based climate model development harnessing machine learning: I. A calibration tool for parameterization improvement, *Journal of Advances in Modeling Earth Systems*, 13, e2020MS002 217, 2021.
- 855 Daleu, C. L., Plant, R., Woolnough, S. J., Sessions, S., Herman, M., Sobel, A., Wang, S., Kim, D., Cheng, A., Bellon, G., et al.: Intercomparison of methods of coupling between convection and large-scale circulation: 1. Comparison over uniform surface conditions, *Journal of Advances in Modeling Earth Systems*, 7, 1576–1601, 2015.
- Del Genio, A. D., Wu, J., Wolf, A. B., Chen, Y., Yao, M.-S., and Kim, D.: Constraints on cumulus parameterization from simulations of observed MJO events, *Journal of Climate*, 28, 6419–6442, 2015.
- 860 Diallo, F., Hourdin, F., Rio, C., Traore, A.-K., Mellul, L., Guichard, F., and Kergoat, L.: The surface energy budget computed at the grid-scale of a climate model challenged by station data in West Africa, *Journal of Advances in Modeling Earth Systems*, 9, 2710–2738, 2017.
- Dirmeyer, P. A., Cash, B. A., Kinter, J. L., Jung, T., Marx, L., Satoh, M., Stan, C., Tomita, H., Towers, P., Wedi, N., et al.: Simulating the diurnal cycle of rainfall in global climate models: Resolution versus parameterization, *Climate dynamics*, 39, 399–418, 2012.
- Dorrestijn, J., Crommelin, D. T., Siebesma, A. P., and Jonker, H. J.: Stochastic parameterization of shallow cumulus convection estimated 865 from high-resolution model data, *Theoretical and Computational Fluid Dynamics*, 27, 133–148, 2013.
- Droegemeier, K. K. and Wilhelmson, R. B.: Numerical simulation of thunderstorm outflow dynamics. Part I: Outflow sensitivity experiments and turbulence dynamics, *Journal of atmospheric sciences*, 44, 1180–1210, 1987.
- Emanuel, K. A.: A scheme for representing cumulus convection in large-scale models, *Journal of Atmospheric Sciences*, 48, 2313–2329, 1991.
- 870 Feng, Z., Hagos, S., Rowe, A. K., Burleyson, C. D., Martini, M. N., and de Szoeke, S. P.: Mechanisms of convective cloud organization by cold pools over tropical warm ocean during the AMIE/DYNAMO field campaign, *Journal of Advances in Modeling Earth Systems*, 7, 357–381, 2015.

- Freitas, S. R., Grell, G. A., Chovert, A. D., Silva Dias, M. A. F., and de Lima Nascimento, E.: A parameterization for cloud organization and propagation by evaporation-driven cold pool edges, *Journal of Advances in Modeling Earth Systems*, 16, e2023MS003 982, 2024.
- 875 Gettelman, A., Truesdale, J., Bacmeister, J., Caldwell, P., Neale, R., Bogenschutz, P., and Simpson, I.: The Single Column Atmosphere Model version 6 (SCAM6): Not a scam but a tool for model evaluation and development, *Journal of Advances in Modeling Earth Systems*, 11, 1381–1401, 2019.
- Grandpeix, J.-Y. and Lafore, J.-P.: A density current parameterization coupled with Emanuel’s convection scheme. Part I: The models, *Journal of the Atmospheric Sciences*, 67, 881–897, 2010.
- 880 Grandpeix, J.-Y., Phillips, V., and Tailleux, R.: Improved mixing representation in Emanuel’s convection scheme, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 130, 3207–3222, 2004.
- Grandpeix, J.-Y., Lafore, J.-P., and Cheruy, F.: A density current parameterization coupled with Emanuel’s convection scheme. Part II: 1D simulations, *Journal of the Atmospheric Sciences*, 67, 898–922, 2010.
- 885 Grant, L. D., Lane, T. P., and van den Heever, S. C.: The role of cold pools in tropical oceanic convective systems, *Journal of the Atmospheric Sciences*, 75, 2615–2634, 2018.
- Guichard, F., Petch, J., Redelsperger, J.-L., Bechtold, P., Chaboureau, J.-P., Cheinet, S., Grabowski, W., Grenier, H., Jones, C., Köhler, M., et al.: Modelling the diurnal cycle of deep precipitating convection over land with cloud-resolving models and single-column models, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical*
- 890 *oceanography*, 130, 3139–3172, 2004.
- Haerter, J. O. and Schlemmer, L.: Intensified cold pool dynamics under stronger surface heating, *Geophysical Research Letters*, 45, 6299–6310, 2018.
- Hourdin, F., Couvreux, F., and Menut, L.: Parameterisation of the dry convective boundary layer based on a mass flux representation of thermals, *Journal of the Atmospheric Sciences*, 59, 1105–1123, 2002.
- 895 Hourdin, F., Musat, I., Bony, S., Braconnot, P., Codron, F., Dufresne, J.-L., Fairhead, L., Filiberti, M.-A., Friedlingstein, P., Grandpeix, J.-Y., et al.: The LMDZ4 general circulation model: climate performance and sensitivity to parametrized physics with emphasis on tropical convection, *Climate Dynamics*, 27, 787–813, 2006.
- Hourdin, F., Foujols, M.-A., Codron, F., Guemas, V., Dufresne, J.-L., Bony, S., Denvil, S., Guez, L., Lott, F., Ghattas, J., et al.: Impact of the LMDZ atmospheric grid configuration on the climate and sensitivity of the IPSL-CM5A coupled model, *Climate Dynamics*, 40,
- 900 2167–2192, 2013.
- Hourdin, F., Jam, A., Rio, C., Couvreux, F., Sandu, I., Lefebvre, M.-P., Briant, F., and Idelkadi, A.: Unified Parameterization of Convective Boundary Layer Transport and Clouds With the Thermal Plume Model, <https://doi.org/10.1029/2019MS001666>, 2019.
- Hourdin, F., Rio, C., Grandpeix, J.-Y., Madeleine, J.-B., Cheruy, F., Rochetin, N., Jam, A., Musat, I., Idelkadi, A., Fairhead, L., et al.: LMDZ6A: The atmospheric component of the IPSL climate model with improved and better tuned physics, *Journal of Advances in*
- 905 *Modeling Earth Systems*, 12, e2019MS001 892, 2020.
- Hourdin, F., Rio, C., Jam, A., Traore, A.-K., and Musat, I.: Convective Boundary Layer Control of the Sea Surface Temperature in the Tropics, *J. of Adv. in Modeling Earth Systems*, 12, e01988, <https://doi.org/10.1029/2019MS001988>, 2020.
- Hourdin, F., Williamson, D., Rio, C., Couvreux, F., Roehrig, R., Villefranche, N., Musat, I., Fairhead, L., Diallo, F. B., and Volodina, V.: Process-based climate model development harnessing machine learning: II. Model calibration from single column to global, *Journal of*
- 910 *Advances in Modeling Earth Systems*, 13, e2020MS002 225, 2021.

- Jam, A., Hourdin, F., Rio, C., and Couvreur, F.: Resolved Versus Parametrized Boundary-Layer Plumes. Part III: Derivation of a Statistical Scheme for Cumulus Clouds, *Boundary Layer Meteorology*, 147, 421–441, <https://doi.org/10.1007/s10546-012-9789-3>, 2013.
- Jeevanjee, N. and Romps, D. M.: Convective self-aggregation, cold pools, and domain size, *Geophysical Research Letters*, 40, 994–998, 2013.
- 915 Kendon, E. J., Roberts, N. M., Senior, C. A., and Roberts, M. J.: Realism of rainfall in a very high-resolution regional climate model, *Journal of Climate*, 25, 5791–5806, 2012.
- Khairoutdinov, M. and Randall, D.: High-resolution simulation of shallow-to-deep convection transition over land, *Journal of the atmospheric sciences*, 63, 3421–3436, 2006.
- Khairoutdinov, M. F. and Randall, D. A.: Cloud resolving modeling of the ARM summer 1997 IOP: Model formulation, results, uncertainties, and sensitivities, *Journal of the Atmospheric Sciences*, 60, 607–625, 2003.
- 920 Krueger, S. K., Morrison, H., and Fridlind, A. M.: Cloud-resolving modeling: ARM and the GCSS story, *Meteorological Monographs*, 57, 25–1, 2016.
- Kurowski, M. J., Suselj, K., Grabowski, W. W., and Teixeira, J.: Shallow-to-deep transition of continental moist convection: Cold pools, surface fluxes, and mesoscale organization, *Journal of the Atmospheric Sciences*, 75, 4071–4090, 2018.
- 925 Lac, C., Chaboureau, J.-P., Masson, V., Pinty, J.-P., Tulet, P., Escobar, J., Leriche, M., Barthe, C., Aouizerats, B., Augros, C., et al.: Overview of the Meso-NH model version 5.4 and its applications, *Geoscientific Model Development*, 11, 1929–1969, 2018.
- Lafore, J.-P. and Moncrieff, M. W.: A numerical investigation of the organization and interaction of the convective and stratiform regions of tropical squall lines, *Journal of Atmospheric Sciences*, 46, 521–544, 1989.
- Legay, A., Deremble, B., and Burchard, H.: Derivation and implementation of a non-local term to improve the oceanic convection representation within the $k-\epsilon$ parameterization, *Journal of Advances in Modeling Earth Systems*, 17, e2024MS004 243, 2025.
- 930 Lochbihler, K., Lenderink, G., and Siebesma, A. P.: Cold pool dynamics shape the response of extreme rainfall events to climate change, *Journal of Advances in Modeling Earth Systems*, 13, e2020MS002 306, 2021.
- Lothon, M., Campistron, B., Chong, M., Couvreur, F., Guichard, F., Rio, C., and Williams, E.: Life cycle of a mesoscale circular gust front observed by a C-band Doppler radar in West Africa, *Monthly weather review*, 139, 1370–1388, 2011.
- 935 Ma, H.-Y., Chuang, C., Klein, S., Lo, M.-H., Zhang, Y., Xie, S., Zheng, X., Ma, P.-L., Zhang, Y., and Phillips, T.: An improved hindcast approach for evaluation and diagnosis of physical processes in global climate models, *Journal of Advances in Modeling Earth Systems*, 7, 1810–1827, 2015.
- Meyer, B. and Haerter, J. O.: Mechanical forcing of convection by cold pools: Collisions and energy scaling, *Journal of Advances in Modeling Earth Systems*, 12, e2020MS002 281, 2020.
- 940 Pantillon, F., Knippertz, P., Marsham, J. H., and Birch, C. E.: A parameterization of convective dust storms for models with mass-flux convection schemes, *Journal of the Atmospheric Sciences*, 72, 2545–2561, 2015.
- Park, S.: A unified convection scheme (UNICON). Part I: Formulation, *Journal of the Atmospheric Sciences*, 71, 3902–3930, 2014.
- Park, S., Song, C., Kim, S., and Kim, J.: Parameterization of the elevated convection with a unified convection scheme (UNICON) and its impacts on the diurnal cycle of precipitation, *Journal of Advances in Modeling Earth Systems*, 16, e2023MS003 651, 2024.
- 945 Provod, M., Marsham, J., Parker, D., and Birch, C.: A characterization of cold pools in the West African Sahel, *Monthly Weather Review*, 144, 1923–1934, 2016.
- Qian, L., Young, G. S., and Frank, W. M.: A convective wake parameterization scheme for use in general circulation models, *Monthly weather review*, 126, 456–469, 1998.

- Randall, D., Khairoutdinov, M., Arakawa, A., and Grabowski, W.: Breaking the cloud parameterization deadlock, *Bulletin of the American Meteorological Society*, 84, 1547–1564, 2003.
- 950 Redelsperger, J.-L., Diedhiou, A., Flamant, C., Janicot, S., Lafore, J.-P., Lebel, T., Polcher, J., Bourlès, B., Caniaux, G., de Rosnay, P., et al.: Amma, une étude multidisciplinaire de la mousson ouest-africaine, *La meteorologie*, 54, 22–32, 2006.
- Rio, C. and Hourdin, F.: A thermal plume model for the convective boundary layer: Representation of cumulus clouds, *Journal of the atmospheric sciences*, 65, 407–425, 2008.
- 955 Rio, C., Hourdin, F., Grandpeix, J.-Y., and Lafore, J.-P.: Shifting the diurnal cycle of parameterized deep convection over land, *Geophysical Research Letters*, 36, 2009.
- Rio, C., Grandpeix, J.-Y., Hourdin, F., Guichard, F., Couvreux, F., Lafore, J.-P., Fridlind, A., Mrowiec, A., Roehrig, R., Rochetin, N., et al.: Control of deep convection by sub-cloud lifting processes: the ALP closure in the LMDZ5B general circulation model, *Climate dynamics*, 40, 2271–2292, 2013.
- 960 Rochetin, N., Grandpeix, J.-Y., Rio, C., and Couvreux, F.: Deep convection triggering by boundary layer thermals. Part II: Stochastic triggering parameterization for the LMDZ GCM, *Journal of the Atmospheric Sciences*, 71, 515–538, 2014.
- Rochetin, N., Hohenegger, C., Touzé-Peiffer, L., and Villefranque, N.: A physically based definition of convectively generated density currents: Detection and characterization in convection-permitting simulations, *Journal of Advances in Modeling Earth Systems*, 13, e2020MS002402, 2021.
- 965 Rooney, G. G., Stirling, A. J., Stratton, R. A., and Whittall, M.: C-POOL: A scheme for modelling convective cold pools in the Met Office Unified Model, *Quarterly Journal of the Royal Meteorological Society*, 148, 962–980, 2022.
- Rotunno, R., Klemp, J. B., and Weisman, M. L.: A theory for strong, long-lived squall lines, *Journal of Atmospheric Sciences*, 45, 463–485, 1988.
- Sadourny, R.: January and July performances of the LMD general circulation model, *New perspectives in climate modeling*, 1984.
- 970 Siebesma, A. P., Bretherton, C. S., Brown, A., Chlond, A., Cuxart, J., Duynkerke, P. G., Jiang, H., Khairoutdinov, M., Lewellen, D., Moeng, C.-H., et al.: A large eddy simulation intercomparison study of shallow cumulus convection, *Journal of the Atmospheric Sciences*, 60, 1201–1219, 2003.
- Stephens, G. L., L’Ecuyer, T., Forbes, R., Gettelmen, A., Golaz, J.-C., Bodas-Salcedo, A., Suzuki, K., Gabriel, P., and Haynes, J.: Dreary state of precipitation in global models, *Journal of Geophysical Research: Atmospheres*, 115, 2010.
- 975 Strauss, C., Ricard, D., Lac, C., and Verrelle, A.: Evaluation of turbulence parametrizations in convective clouds and their environment based on a large-eddy simulation, *Quarterly Journal of the Royal Meteorological Society*, 145, 3195–3217, 2019.
- Suselj, K., Kurowski, M. J., and Teixeira, J.: A unified eddy-diffusivity/mass-flux approach for modeling atmospheric convection, *Journal of the Atmospheric Sciences*, 76, 2505–2537, 2019.
- Tan, J., Oreopoulos, L., Jakob, C., and Jin, D.: Evaluating rainfall errors in global climate models through cloud regimes, *Climate Dynamics*, 50, 3301–3314, 2018.
- 980 Torri, G. and Kuang, Z.: On cold pool collisions in tropical boundary layers, *Geophysical Research Letters*, 46, 399–407, 2019.
- Touzé-Peiffer, L., Vogel, R., and Rochetin, N.: Cold pools observed during eurec 4 a: Detection and characterization from atmospheric soundings, *Journal of Applied Meteorology and Climatology*, 61, 593–610, 2022.
- Vogel, R., Konow, H., Schulz, H., and Zuidema, P.: A climatology of trade-wind cumulus cold pools and their link to mesoscale cloud organization, *Atmospheric Chemistry and Physics*, 21, 16609–16630, 2021.
- 985

- Weisman, M. L. and Rotunno, R.: “A theory for strong long-lived squall lines” revisited, *Journal of the Atmospheric Sciences*, 61, 361–382, 2004.
- Williamson, D., Goldstein, M., Allison, L., Blaker, A., Challenor, P., Jackson, L., and Yamazaki, K.: History matching for exploring and reducing climate model parameter space using observations and a large perturbed physics ensemble, *Climate dynamics*, 41, 1703–1729, 2013.
- 990 Yamada, T.: Simulations of Nocturnal Drainage Flows by a q^2l Turbulence Closure Model, *J. Atmos. Sci.*, 40, 91–106, 1983.
- Yanai, M., Esbensen, S., and Chu, J.-H.: Determination of bulk properties of tropical cloud clusters from large-scale heat and moisture budgets, *Journal of Atmospheric Sciences*, 30, 611–627, 1973.
- Young, G. S., Perugini, S. M., and Fairall, C.: Convective wakes in the equatorial western Pacific during TOGA, *Monthly Weather Review*, 995 123, 110–123, 1995.
- Zhang, M., Somerville, R. C., and Xie, S.: The SCM concept and creation of ARM forcing datasets, *Meteorological Monographs*, 57, 24–1, 2016.
- Zuidema, P., Torri, G., Muller, C., and Chandra, A.: A survey of precipitation-induced atmospheric cold pools over oceans and their interactions with the larger-scale environment, *Surveys in Geophysics*, 38, 1283–1305, 2017.